



UNIVERSITETET I OSLO  
DET MEDISINSKE FAKULTET



# STUDIES ON SUBFRACTIONS OF FIBRINOGEN

*With special emphasis on fibrinogen quantification, viscosity and inflammation*

**Torstein Jensen, MD**



University of Oslo  
Norway  
2008

© **Torstein Jensen, 2008**

*Series of dissertations submitted to the  
Faculty of Medicine, University of Oslo  
No. 597*

ISBN 978-82-8072-883-8

All rights reserved. No part of this publication may be  
reproduced or transmitted, in any form or by any means, without permission.

Cover: Inger Sandved Anfinsen.  
Printed in Norway: AiT e-dit AS, Oslo, 2008.

Produced in co-operation with Unipub AS.  
The thesis is produced by Unipub AS merely in connection with the  
thesis defence. Kindly direct all inquiries regarding the thesis to the copyright  
holder or the unit which grants the doctorate.

*Unipub AS is owned by  
The University Foundation for Student Life (SiO)*

## CONTENTS

	Page
<b>1. ACKNOWLEDGEMENTS.....</b>	<b>5</b>
<b>2. ABBREVIATIONS.....</b>	<b>6</b>
<b>3. LIST OF PAPERS.....</b>	<b>8</b>
<b>4. GENERAL INTRODUCTION.....</b>	<b>9</b>
4.1 Fibrinogen structure .....	9
4.2 Synthesis.....	12
4.3 Conversion of fibrinogen to fibrin.....	14
4.4 Metabolism of fibrinogen and fibrin.....	18
4.4 Fibrinogen and risk of cardiovascular disease.....	19
4.5 Fibrinogen and plasma viscosity.....	20
4.6 Fibrinogen and acute-phase.....	21
4.7 Fibrinolysis.....	22
4.8 Fibrinogen assays.....	23
4.9 Interaction with blood platelets and leukocytes.....	24
<b>5. AIMS OF THE STUDY.....</b>	<b>25</b>
<b>6. MATERIALS AND METHODS.....</b>	<b>28</b>
<b>7. SUMMARY OF RESULTS.....</b>	<b>34</b>
6.1 Paper I.....	34

<b>6.2 Paper II.....</b>	<b>34</b>
<b>6.3 Paper III.....</b>	<b>35</b>
<b>6.4 Paper IV.....</b>	<b>36</b>
<b>6.5 Paper V.....</b>	<b>36</b>
<b>8. GENERAL DISCUSSION.....</b>	<b>37</b>
<b>9. CONCLUSIONS.....</b>	<b>47</b>
<b>10. REFERENCES.....</b>	<b>48</b>
<b>11. PAPERS I-V</b>	

## 1. ACKNOWLEDGEMENTS

The present work was initiated in 1999, and the laboratory work carried out at the Haematological Research Laboratory, Medical Division, Ullevål University Hospital. I would like to thank my supervisor Professor Ole Henning Skjøsberg, Department of Pulmonary Medicine, for his guidance through these years. I have greatly appreciated our meetings and his valuable and constructive advice and excellent evaluations. His inspiration and encouragement has been of great importance. Always at hand whenever I needed help, yet I have felt free to pursue my own ideas and interests. I am thankful for having the opportunity to work with him.

The cooperation between him and my co-supervisor Professor Emeritus Hans Christian Godal, the founder of Haematological Research Laboratory, was truly excellent. Godal had a life-long interest in fibrinogen research, and I am grateful for his good ideas and enthusiasm in my work. Although long into his retirement, he attended the laboratory almost daily, until he passed away in 2005. Through our regular discussions I learned much about fibrinogen research.

I would like to thank Professor Per Morten Sandset, head of the Department of Haematology and Haematological Research Laboratory, for providing excellent working conditions, and for his extensive and valuable contributions to my work. I am also grateful to the staff at the Haematological Research Laboratory for aiding me in the laboratory work. Special thanks to Karl Gravem, Trine Andersen, Brit Steinsvik and Kaia Haugbro for expert assistance during my experiments, and to Dr Sigrun Halvorsen, Department of Cardiology, for her significant and valuable contributions.

I would like to express my gratitude to Professor Emeritus Peter Kierulf and Dr Olav Klingenberg at the Department of Clinical Chemistry for excellent cooperation, and to Gunn Brit Joø and Reidun Øvstebø for expert technical assistance. Finally, I would like to thank my family and my friends for supporting me through these years. Special thanks to my colleague and good friend Dr. Svein Solheim, Lovisenberg Hospital, for his help and valuable advice.

I dedicate this thesis to the memory of Hans Christian Godal.

Oslo, February 2008

Torstein Jensen

## 2. ABBREVIATIONS

aa	amino acids
$\alpha$ C	carboxy terminal end of the fibrinogen $\alpha$ -chain
ADP	adenosine diphosphate
CHD	coronary heart disease
CRP	C-reactive protein
EDTA	ethylene-diamine-tetra-acetic acid
FgDP	fibrinogen degradation products
FnDP	fibrin degradation products
FPA	fibrinopeptide A
FPB	fibrinopeptide B
HMW	high molecular weight
IHD	ischemic heart disease
IL-1	interleukin 1
IL-6	interleukin 6
ISTH	international society of thrombosis and haemostasis
kDa	kiloDalton
LMW	low molecular weight
LMW <sup>v</sup>	very low molecular weight
LPS	lipopolysaccharid
M	molar
mPa.s	millipascal seconds
MW	molecular weight
N-terminal	amino-terminal
PAGE	polyacrylamid gel electrophoresis
PAI	plasminogen activator inhibitor
PBMC	peripheral blood mononuclear cells
PBS	phosphate buffered saline
RNA	ribonucleic acid
SDS	sodium dodecyl sulphate

SD	standard deviation
TAFI	thrombin-activatable fibrinolysis inhibitor
TF	tissue factor
TNF	tumor necrosis factor
U-PA	urokinase-type plasminogen activator
U	unit
vWF	von Willebrand Factor

### **3. LIST OF PAPERS**

#### **PAPER I**

Influence of freeze-drying on the clotting properties of fibrinogen in plasma.

Torstein Jensen, Sigrun Halvorsen, Hans C. Godal, Ole H. Skjøsberg

Thrombosis Research 2002; 105: 499– 502.

#### **PAPER II**

Discrepancy between fibrinogen concentrations determined by clotting rate and clottability assays during the acute-phase reaction.

Torstein Jensen, Sigrun Halvorsen, Hans C. Godal, Per M. Sandset and Ole H. Skjøsberg. Thrombosis Research 2000; 100: 397-403.

#### **PAPER III**

The viscosity of fibrinogen subfractions and of EDTA denatured fibrinogen do not differ from that of native fibrinogen.

Torstein Jensen, Sigrun Halvorsen, Hans C. Godal, Per M. Sandset, Ole H. Skjøsberg. Thrombosis Research 2004; 113: 51–56.

#### **PAPER IV**

A daily glass of red wine induces a prolonged reduction in plasma viscosity: a randomized controlled trial.

Torstein Jensen, Lars J. Retterstøl, Per M. Sandset, Hans C. Godal, Ole H. Skjøsberg. Blood Coagulation and Fibrinolysis 2006; 17: 471–476.

#### **PAPER V**

Fibrinogen and fibrin induce synthesis of proinflammatory cytokines from isolated peripheral blood mononuclear cells.

Torstein Jensen, Peter Kierulf, Olav Klingenberg, Gunn Brit Joø, Hans C. Godal, Per M. Sandset, Ole H. Skjøsberg.

Thrombosis and Haemostasis 2007; 20: 822-829.



## 4. GENERAL INTRODUCTION

Fibrinogen is a large, soluble glycoprotein. It plays a central role in human hemostasis through its conversion to fibrin in the last step of the coagulation cascade, and by its essential role in platelet aggregation. Normal plasma concentration is 1.7-4.0 g/l. Being an acute-phase protein, the concentration rises rapidly during inflammatory conditions, major surgery and certain cancer forms (1-3). The plasma concentration results from a complex interaction between genetic and environmental factors (4). Plasma fibrinogen levels are higher in females than in males, and increased concentrations are found in postmenopausal women, smokers, patients with diabetes, obesity, hypercholesterolemia and hypertension (5). Factors which may decrease fibrinogen levels include physical exercise, weight loss, moderate alcohol consumption and smoking cessation (4;6-10). Treatment with fibrates and platelet aggregation inhibiting drugs such as ticlopidine may lower plasma levels of fibrinogen (11). Different responses of hormone replacement therapy have been reported (9;12-14). Genetic heritability accounts for about 50% of the variance in fibrinogen concentration between individuals (15).

### Structure

In 1959 Hall and Slayter produced electron microscopy images of fibrinogen revealing a rod-like molecule with 3 interconnected globules (16). The length was estimated to be  $475 \pm 25$  Å. The diameter of the center globule was approximately 50 Å, and the distal globules were approximately 65 Å (16;17). The structure was later confirmed by other investigators (18;19). Using atomic force microscopy, Marchant and colleagues obtained three dimensional images of fibrinogen molecules under aqueous conditions, showing a mixture of monomers, dimers and trimeric structures linked through D-D globular interconnections (20).

The fibrinogen molecule is a dimer consisting of 3 pairs of the non-identical polypeptide chains A $\alpha$ , B $\beta$  and  $\gamma$  with molecular weight 340 kDa (21). The nomenclature (A $\alpha$ ,B $\beta$ , $\gamma$ )<sub>2</sub> reflects the peptides A and B cleaved from fibrinogen by thrombin and the longer  $\alpha$ -,  $\beta$ - and  $\gamma$ -chains. No peptides are split from the  $\gamma$ -chain during the conversion of fibrinogen

to fibrin. The total number of amino-acids is 2964, and the complete amino-acid




Figure 1. Electron microscope images showing fibrinogen molecules with a central (E)- and two peripheral (D) regions (1A), fibrin protofibrils with half-staggered overlapping fibrin monomers (1B), and thicker, branching fibrin fibers (1C). Reproduced from Weisel (22).

sequence was presented in 1980 by Henschen and Lottspeich (23;24). The molecular weights of the individual chains are 66.5 kDa, (610 amino-acids), 52 kDa (461 amino-acids), and 46.5 kDa (411 amino-acids) for the A $\alpha$ -, B $\beta$ - and  $\gamma$ -chain, respectively (25;26). The 6 polypeptide chains are gathered in the central E region (E-domain) (27), which is connected to the 2 D regions (D-domains) of the molecule by the “interconnector”, comprising 111-112 amino acids of the A $\alpha$ -, B $\beta$ -, and  $\gamma$ -chains joined together in a coiled-coil (28) (figure 2). The D region can be further divided into a distal nodule containing the C-terminal part of the  $\gamma$ -chain ( $\gamma$ C), and the proximal nodule including the C-terminal B $\beta$ -chain. The C-terminal part of the A $\alpha$ -chain ( $\alpha$ C) consists of an elongated polypeptide extending from the D region, interacting with the central E region, and forming intra-molecular  $\alpha$ C- $\alpha$ C connections (29;30).




Figure 2. Schematic drawing of the fibrinogen molecule showing the N- and C-terminal ends of the A $\alpha$ -, B $\beta$ -, and  $\gamma$ -chains. The coiled-coil regions are illustrated with diagonal lines, and disulfide bonds marked with solid lines. CHO indicates carbohydrate attachment sites, and P and T represent major cleavage sites for plasmin and thrombin, respectively. Adapted from Hantgan (31).

### **Calcium**

Calcium is important for the structural integrity and function of the fibrinogen molecule. High affinity binding sites for calcium are located in the  $\gamma$ -chain at Asp<sup>318</sup>, Asp<sup>320</sup>, Gly<sup>324</sup> and Phe<sup>322</sup>. In addition several low affinity calcium binding sites exists which are less well defined (32). In plasma the binding sites are usually occupied providing protection from enzymatic degradation (33;34).

### **Disulphide bonds**

Twenty nine disulfide chains located in the A $\alpha$ - (8), B $\beta$ - (11) and  $\gamma$ -chains (10) hold the 6 chains together (35). The central E-region contains the 2 pairs of A and B peptides linked together by three inter-chained disulfide bonds, 1 between the A $\alpha$ -chains and 2 between the  $\gamma$ -chains. One disulfide bridge links the A $\alpha$ - and the B $\beta$ -chain.

### **Carbohydrate chains**

Fibrinogen is a glycoprotein containing four oligosaccharides located at B $\beta$ <sup>364</sup> and  $\gamma$ <sup>52</sup> and linked through N-glycosyl bonds. The  $\alpha$ -chain contains no carbohydrate side chains (36). The carbohydrate chains are of major importance for the solubility of fibrinogen, and modifications of the carbohydrate chains affect the fibrin polymerization properties and clot structure. Fibrinogen contains equal amounts of mono- and disialicated chains, and variations of the sialic-acid content of the carbohydrate chains account for some of the

heterogeneity of fibrinogen in plasma. Complete deglycosylation results in accelerated fibrin polymerization, increased lateral aggregation with thicker, less branched fibrin fibers and a more porous network (37).

### **Synthesis of fibrinogen**

Fibrinogen in plasma is mainly produced in the liver (38;39) with a half-life in plasma of 3-5 days (40-42). Synthesis under normal conditions range from 1.7-5 g/day (43). Fibrinogen is an acute-phase protein, and during acute-phase conditions such as major surgery, bacterial infections and trauma, gene transcription and release of fibrinogen rises rapidly. Cytokines in the interleukin-6 (IL-6) family have been recognized as important stimulants to increased fibrinogen synthesis. Plasma fibrinogen levels can, thus, increase manifold. Fibrinogen and fibrin degradation products (FgDP/FnDP) have been shown to stimulate fibrinogen synthesis in vitro (44;45). The results are inconsistent, however, and the physiological role of FgDP and FnDP on fibrinogen synthesis in vivo is not clear (44-48). Although 75% of the total fibrinogen is found in plasma, it may also be found in lymph nodes, blood platelets and in interstitial fluid. In vitro studies have shown that fibrinogen may also be synthesized by non-hepatic cell lines such as epithelial cells from lung and intestine as well as trophoblasts and granulosa cells (49-52). The physiological role of this locally produced fibrinogen is not clear. Although the A $\alpha$ , B $\beta$ , and  $\gamma$  chains are encoded by dedicated genes closely linked on the distal third of the long arm of chromosome 4, bands q23-32 (53), the transcription of the 3 different genes are closely coordinated (54). The gene for the A $\alpha$ -chain, located at the centre of the fibrinogen cluster, consists of 5 exons that translate into 625 amino acids, 15 of which are removed after transcription. The gene coding for the B $\beta$ -chains contain 8 exons and codes for a 461 residue polypeptide and the gene coding for  $\gamma$ -chain comprise 10 exons coding for the 411 residue  $\gamma$ -chain (53).

The intracellular assembly of the 3 chains in cytoplasmatic reticulum joins single chains to a 3-chained half-molecule which is subsequently joined to the final dimeric fibrinogen molecule. When hepatocytes are stimulated with IL-6, the genes for the 3 subunits are upregulated simultaneously and to the same extent (55;56). Cytokines such as

interleukin-1 $\beta$  and tumor necrosis factor- $\alpha$  and glucocorticosteroids may modify the IL-6 response (54;57).

### **Variations of the $\alpha$ -chain**

Heterogeneity in circulating fibrinogen molecules are dominated by heterogeneity of the C-terminal  $\alpha$ -chain (58;59). Depending on the degree of C-terminal  $\alpha$ -chain degradation, 3 main fibrinogen fractions have been described: High molecular weight (HMW-) fibrinogen (MW 340 kDa) having both its  $\alpha$ -chains intact; low molecular weight (LMW-) fibrinogen (305 kDa), which has lost the C-terminal end of one  $\alpha$ -chain, and very low molecular weight (LMW'-) fibrinogen (280 kDa), which has lost the C-terminal end of both  $\alpha$ -chains (2;60). These fractions constitute approximately 70%, 25% and 5%, respectively, of the total plasma fibrinogen content. Experiments using western-blotting and immunostaining techniques have, however, revealed a considerable larger diversity of A $\alpha$ -chain remnants with intact N-terminal ends and MW 66,200 to 36,000 Da in normal plasma (61). High molecular weight fibrinogen has lower solubility, shorter thrombin clotting time and increased clottability compared to the low molecular forms, illustrating the importance of the carboxy- terminal  $\alpha$ -chain for the clotting properties of fibrinogen (62-68). Furthermore, clots from  $\alpha$ C-degraded fibrinogen have decreased turbidity and lack mechanical strength compared to native fibrin clots. Although not essential for clot formation, the loss of one or both  $\alpha$ C-terminals significantly retards the polymerization process (29;69). In 1992 an extended version of the  $\alpha$ -chain,  $\alpha$ E, was described (70;71). Based on predicted mass, fibrinogen molecules containing the 236 residue extended  $\alpha$ -chains are called Fibrinogen 420 (MW 420 kDa), to distinguish it from the more predominant Fibrinogen 340. Fibrinogen 420 comprises approximately 1 % of the total fibrinogen concentration in adults (72). It has been shown to be involved in leukocyte adhesion and migration through binding of the  $\alpha$ <sub>E</sub>C terminal to the leukocyte integrins  $\alpha$ <sub>M</sub> $\beta$ 2 and  $\alpha$ <sub>X</sub> $\beta$ 2 (73).

### **Variations of the $\gamma$ -chain**

Alternative processing of the  $\gamma$ -chain gene transcript gives rise to an alternative  $\gamma$ -chain,  $\gamma'$ , with a negatively charged 20 amino acid extension at the C terminal end of the  $\gamma$ -chain, substituting the  $\gamma$  408–411 with a 20 amino acid sequence,  $\gamma'$ 408–427, that contains two sulfated tyrosines. The  $\gamma'$ -chain constitutes about 8% of the  $\gamma$  chains (74–76), and in plasma 10–15% of the fibrinogen molecules contain the  $\gamma'$  variant (77). The  $\gamma'$ -chain includes binding sites for thrombin and the B chain of factor XIII, suggesting that the  $\gamma'$ -chain function as a carrier for factor XIII in the circulation, making it readily accessible during fibrin formation (78). The  $\gamma$ - and  $\gamma'$ -chains are equally effective cross linked by XIIIa. Due to lack of the full platelet binding sequence 400–411, the  $\gamma'$ -chain does not support ADP induced fibrinogen mediated platelet aggregation (79;80).

### **Conversion of fibrinogen to fibrin**

The conversion of soluble fibrinogen to an insoluble fibrin network is initiated by the thrombin mediated removal of the fibrinopeptides A and B from the N-terminal A $\alpha$ - and B $\beta$ -chains (81). The cleavage of FPA at amino acids 15–16 occurs more rapidly than the split of FPB between amino acids 16–17 (82), which has been suggested to be involved in lateral growth and branching of the fibrin polymer (83). Thus, 2 FPA and 2 FPB are released from each fibrinogen molecule exposing the peptide sequence Gly-His-Arg-Pro-amide in the  $\alpha$ -knob, and the Gly-Pro-Arg-Pro-amide from the  $\beta$ -knob. Complimentary “holes” are located in the globular C-terminal  $\beta$ - and  $\gamma$ -chains. Although structurally similar, the  $\beta$ C- and  $\gamma$ C-holes are able to distinguish between these peptides.

### *Thrombin*

Thrombin is a serine protease produced in the liver as the 70 kD zymogene prothrombin (84). The 579 residue prothrombin undergoes posttranscriptional modifications before it is secreted into the circulation. Following damage to the blood vessel and exposition of subendothelial tissue factor (TF), factor X is converted to Xa in the presence of FVIIa and  $\text{Ca}^{2+}$  in the initial phase of coagulation. Prothrombin is subsequently converted to the 35 kDa biologically active thrombin by the prothrombinase complex, including FVa, FXa, and anionic phospholipids in the presence of  $\text{Ca}^{2+}$ . Thus, prothrombin is cleaved

after Arg<sup>320</sup> forming the intermediate mezothrombin and subsequently after Arg<sup>271</sup> to form thrombin (85). Thrombin comprises a 49 residue light chain and a 259 residue heavy chain linked together by a single disulphide bridge. The heavy chain is intra-connected through 3 disulphide bonds. The initial small amounts of thrombin formed by the extrinsic pathway trigger the intrinsic coagulation pathway through activation of FXI, FVIII and FV, thus, creating more thrombin.

The activity half-life of thrombin is only 14 seconds, and it is rapidly inactivated by inhibitors. The principal circulating thrombin inhibitors are antithrombin, heparin cofactor II (HCII) and protease nexin 1 (86;87). However, thrombin also has the ability to inhibit its own generation through interaction with endothelial bound thrombomodulin, which activates the protein C pathway. Protein C and protein S effectively inhibit factors Va and VIIIa, thus preventing further thrombin generation. The proteolytic properties of thrombin are affected through interaction with Na<sup>+</sup>. Several Na<sup>+</sup> binding sites have been identified, the principal binding sites being confined to the C-terminal region of the heavy chain (88;89). Binding of Na<sup>+</sup> to thrombin alters the protein structure and allosterically enhances its catalytic properties. At physiological Na<sup>+</sup> concentrations, the sodium binding sites in thrombin are fully occupied.

### *Polymerization*

The removal of the acidic FPA from fibrinogen changes the N-terminal net charge, and is followed by a spontaneous polymerization step in which two fibrin monomers (des-AA fibrin) are arranged in a non-covalent bonded staggered overlap between the E-domain and complimentary binding sites in the C-terminal  $\gamma$ -chain of the D-domain (90). In addition to the E-/D-domain interaction, D/D-domain intermolecular bonds form in the fibrin oligomer, thus forming a double stranded protofibril. The release of fibrinopeptide B, although starting at the same time as the release of FPA, occurs at a slower rate than the release of FPA (82;91). Fibrin gel formation, however, accelerates the FPB release rate about sevenfold (92;93). The removal of FPB exposes an additional polymerization site in the E-domain called B “knobs” which are complementary to b “holes” in the C-terminal  $\beta$ -chain (94;95). Removal of the two FPBs induces a conformational change of the  $\alpha$ -chain. The two  $\alpha$ -chains, usually stretching parallel to the coiled coil and meeting in

the central E-domain, dissociate and form a mobile extension from the D-domain. This allows interaction between neighboring  $\alpha$ -chains, which promotes lateral aggregation of fibrin protofibrils and strengthening of the mechanical properties of the clots (96). The non-covalent ligation between the E-/D-domains are easily broken by solvents such as urea, acetic acid and sodium bromide. Active “knobs” in fibrin monomers or fibrin protofibrils may bind to always exposed “holes” in circulating fibrinogen forming soluble fibrinogen-fibrin complexes (97;98), thus, inhibiting fibrin polymerization. Even degradation products of fibrin and fibrinogen may interact with “holes” in the D-domain and affect clot formation

The fibrin clot architecture is influenced by several factors. High fibrinogen levels and fast activation leads to thin fibrin fibers and more rigid clots with smaller pores, making it less susceptible to fibrinolytics. Slow activation, on the other hand, leads to thicker fibrin strands with larger pores (99). Ionic strength, increased glucose or homocysteine levels, the presence of other plasma proteins (100) and cross linking of proteins such as PAI-2, TAFI,  $\alpha$ -2 antiplasmin as well as fibrinogen- and factor XIII-polymorphisms may all affect the fibrin clot structure (for review see (101)).






Figure 3. Illustration of the conversion of fibrinogen to fibrin by thrombin, fibrin polymerization and factor XIIIa mediated cross-linking according to Weisel (22).

### *Cross-linking*

The fibrin matrix formed by the action of thrombin is fragile. Through factor XIIIa mediated cross-linking of the  $\gamma$ -chains the fibrin clot is strengthened and made resistant to physical stress and proteolytic attacks. Factor XIII in plasma is a 326 kDa tetramer including 2 A chains and 2 B chains. About half of the total factor XIII in blood is contained in platelets in the form of A2 dimers. In plasma, most of the factor XIII zymogen A2B2 is bound to fibrinogen, especially to the  $\gamma$ '-chain of Fibrinogen 420 (102). Factor XIII is activated through the thrombin mediated proteolysis of the A chain. Activated factor XIII induces inter-chain covalent cross-linking between one C-terminal glutamine ( $\gamma$ Gln398/399) and one lysine ( $\gamma$ Lys406) residue in adjacent  $\gamma$ -chains (103;104). Due to the close proximity of the acceptor and donor sites in the fibrin clot,

factor XIIIa catalyses cross-linking of fibrin much faster than cross-linking of fibrinogen (105). Intermolecular  $\alpha$ -chain cross-linking generating  $\alpha$ -oligomers and polymers occur at a slower rate and facilitate lateral fibril association and fiber growth. Crosslinking may also occur between fibrin  $\alpha$ - and  $\gamma$ -chains, and even between fibrinogen  $\gamma$ -chains.

### **Metabolism of fibrinogen and fibrin**

Only 2-3% of the fibrinogen molecules are removed from the circulation as a consequence of coagulation and subsequent fibrinolysis (106). Although plasmin mediated proteolysis is the dominating fibrin degradation pathway, plasmin does not play an important role in the degradation of fibrinogen, except during exogenous treatment with plasminogen activators (107;108). Furthermore, it seems that HMW-fibrinogen only partially is metabolized to LMW-fibrinogen in vivo (108-110). To what extent HMW-fibrinogen is metabolized to LMW-fibrinogen, and which additional mechanisms are involved in the degradation of fibrinogen in vivo are not clear.

In vitro studies have shown that the A $\alpha$ -, B $\beta$ -, and  $\gamma$ -chains differ in their susceptibility to plasmin (111). The A $\alpha$ -chain, with its elongated C-terminal end, is more vulnerable to plasmin attacks than the B $\beta$ - and  $\gamma$ -chains. Similarly, the B $\beta$ -chains are more susceptible to plasmin degradation than the  $\gamma$ -chains. The destruction of the  $\alpha$ C-domain results in the formation of fragment X (MW 250 kDa) and degradation products from the  $\alpha$ -chain named A, B and C (111;112). Further digestion of the coiled-coil of the fragment X, gives rise to fragments D (MW 83-100 kDa) and Y (155 kDa). Fragment Y is quickly degraded to the terminal fragments D (81 kDa) and E (41 kDa) (figure 4). Fragment E corresponds to the N-terminal domain of the native fibrinogen molecule. Fragments D and E constitute the final plasmin mediated fibrinogen degradation products. The affinity of plasminogen to polymerizing fibrin is stronger than to circulating fibrinogen. Although fibrinogen has high affinity binding sites for plasminogen at the C-terminal  $\alpha$ -chain, the formation of the ternary complex fibrin-plasminogen-t-PA is required to accomplish effective t-PA mediated plasminogen activation. The fibrinolysis of cross-linked fibrin creates distinct degradation products such as D-D (D-dimer), comprising two fragments D joined by  $\gamma$ - $\gamma$  cross-links (113;114), and the non-covalent combination of fragment D-D with fragment E (115). Further degradation of D-dimer, and digestion of the C-

terminal end of the A $\alpha$ -chain by human neutrophil elastase has been shown in vitro (64;116).




Figure 4. Illustration of the plasmin mediated degradation of fibrinogen to the intermediary X and Y molecules, and the final degradation products D and E. Reproduced from Gaffney (113).

### Fibrinogen and risk of cardiovascular disease

A number of clinical trials have shown an association between high fibrinogen levels and increased risk of cardiovascular disease (11;117-120). In a metaanalysis comprising 18 prospective studies and 4000 patients with coronary heart disease (CHD), the relative risk of CHD was 1.8 (95 % confidence interval 1.6-2.0) in patients with fibrinogen concentration in the highest tertile compared to the lowest (121). Increased plasma fibrinogen concentrations have also been related to adverse prognosis following myocardial infarction (122;123), increased risk of stroke (124) and increased cardiovascular- and total mortality in asymptomatic patients (118). The observation that fibrinogen is a strong and independent marker for CHD has raised the question whether

high fibrinogen levels promote the atherothrombotic process, or is only a marker of the chronic low-grade inflammation associated with atherosclerotic vascular disease (125).

Possible mechanisms by which fibrinogen may increase CHD risk include its contribution to plasma- (126) and blood viscosity (127;128). Furthermore, a high fibrinogen concentration has been shown to correlate with increased size, weight, and fibrin content of thrombi (129). The key role of fibrinogen in platelet aggregation, and its modulating effect on endothelial function and smooth muscle proliferation, might also affect the atherothrombotic process. On the other hand, studies using genetically modified mice have indicated that fibrinogen is not strictly required for the development of advanced atherosclerotic disease (130).

### **Fibrinogen and plasma viscosity**

Viscosity ( $\eta$ ) can be defined as a fluid's resistance to flow attributable to internal friction between molecules or larger particles (131). Isaac Newton (1642-1727) described a model for the viscous properties of fluids, suggesting that the viscosity is proportional to the velocity at which parts of the fluid are separated from one another due to the flow. Newtonian fluids flow like water, and is characterized by changes in its shear rate being proportional equal to changes in shear stress ( $\gamma = V/H$  where  $V$  is the velocity and  $H$  is the distance between two parallel plates). Shear stress is the applied force per unit area which causes relative movement of fluid layers, and can be described by the equation:

$\tau = \mu \frac{dv}{dx}$  ( $\tau$ = the shear stress exerted by the fluid,  $\mu$ =the fluid viscosity,  $dv/dx$ =the velocity gradient perpendicular to the direction of shear). In non-newtonian fluids, however, the viscosity changes inversely with the shear rate. This implies that for a Newtonian fluid such as plasma, viscosity depends only on pressure, temperature and the chemical composition of the fluid. Different principles have been utilized to determine the viscosity of blood and plasma (132). The "falling-ball" viscometer provides an easy and precise method of measuring the viscosity of Newtonian, low-viscosity fluids such as plasma. By measuring the time taken by a solid sphere to travel a given distance through an inclined glass tube filled with the sample, the viscosity can be calculated.

Albumin, globulins (alpha-1, alpha-2, beta and gamma globulins) and clotting factors constitute the main proteins in plasma. In addition to plasma concentration, the size and shape of proteins determine the effect on plasma viscosity (133). The fibrinogen molecule is a large, rod shaped protein, and fibrinogen has been shown to be a major determinant of plasma viscosity. Variations in plasma viscosity between subjects can partly be attributed to variations in plasma fibrinogen levels (134;135).

### **Fibrinogen and acute-phase**

The initial host response to tissue injury such as infections, major surgery, trauma and some malignant conditions comprises a series of predetermined, coordinated events. This early immune response, called the “acute-phase reaction” (APR), is initiated by the recognition by specific cell receptors of microbial antigens such as lipopolysaccharides (LPS) or other components of the bacterial cell wall (136). The APR includes a local and a systemic response characterized by fever, leukocytosis and increased metabolic rate. In the micro vasculature changes in vascular permeability is observed. Important cytokines involved in the amplification and propagation of the APR include IL-1 $\beta$ , TNF- $\alpha$ , IL-6 and IFN- $\gamma$  (137). Animal studies have indicated a critical role of IL-6 on upregulation of the synthesis and release of acute phase proteins such as clotting-, and complement factors, immune modulating proteins such as fibronectin, CRP, amyloid A and proteins such as haptoglobin, ferritin, ceruloplasmin and  $\alpha_1$ -antitrypsin (137;138). The increased mRNA transcription of the different acute phase proteins is mediated through activation of various nuclear factors (139;140). Plasma fibrinogen concentration may increase 2 to 10 fold during acute phase conditions. Holm and colleagues showed that the increased fibrinogen concentration was mainly due to increased levels of HMW-fibrinogen, with a modest increase in LMW-fibrinogen (2). Due to the increase in HMW-fibrinogen synthesis, the HMW-/LMW-fibrinogen ratio raises 4-8 hours following the initiation of the acute phase, reaching its highest level after 3-4 days (2). The elevated HMW-/LMW-fibrinogen ratio gradually returns to normal levels during the ensuing week, whereas total fibrinogen levels may remain elevated for a longer period of time (2). HMW-fibrinogen has been shown to have increased clotting rate as reflected by shorter thrombin clotting time compared to LMW-, and LMW'-fibrinogen (63). The degree of phosphorylation of

fibrinopeptide A has been shown to increase from 25-30% in normal plasma to more than 50% during acute phase conditions (141).

### **Fibrinolysis**

Activation of coagulation at sites of vascular damage is vital to prevent fatal blood loss. Confining the coagulation process to the place of injury and the subsequent lysis of the clot are, however, equally important to prevent excessive thrombosis and maintain patency of the vasculature. The fibrinolytic system comprises the proenzyme plasminogen, which is converted to the active enzyme plasmin by tissue type plasminogen activator (t-PA) or urokinase type plasminogen activator (u-PA). Plasmin cleaves the fibrin(ogen) molecule at multiple sites resulting in the formation of fibrin(ogen) degradation products (FgDP and FnDP).

### ***Plasminogen***

Plasminogen is a single-chain glycoprotein found in plasma at 1.5-2.0  $\mu\text{M}$  concentrations. The molecule comprises 791 amino acids with MW 92 kDa, including 5 homologous triple-loop structures of MW 10 kDa each (142). The interaction of plasmin with fibrin and  $\alpha_2$ -antiplasmin is mediated by lysine- and aminohexyl binding sites located in the triple-loops. In addition to the native glu-plasminogen, limited plasmic digestion results in modified forms, lys-plasminogen, containing N-terminal lysine, valine or methionine (143). Through cleavage of the Arg<sup>561</sup>-Val<sup>562</sup> peptide bond plasminogen is converted to the active two-chain serine protease plasmin (144). Plasmin is the natural protease for the dissolution of fibrin in vivo.

### ***Plasminogen activators***

Tissue type plasminogen activator (t-PA) is a 527 residue serine protease with MW 70 kDa. Hydrolysis of the Arg<sup>275</sup>-Ile<sup>276</sup> bond converts it to a two chained form, interconnected by a disulfide bond (145). It is released from the vascular endothelium into the vessel lumen as a result of stress, exercise, venous occlusion and specific drugs (146). Circulating t-PA is mostly bound to plasminogen activator inhibitors (PAI) (147). Residues 276-527 constitute the serine protease domain with the active site amino acids

His<sup>322</sup>, Asp<sup>371</sup> and Ser<sup>478</sup> (148). t-PA activity is greatly enhanced by the presence of fibrin, and is further amplified through the subsequent formation of fragment-X through plasmin mediated splitting of the C terminal  $\alpha$ -chain and N-terminal of the  $\beta$ -chain of fibrin (149). In vitro studies suggest that the plasminogen activating capacity of the 3 natural occurring principal fibrin subfractions with different degree of  $\alpha$ -C degradation does not differ (150). Kinetic in vitro studies on the activation of plasminogen in the presence of fibrin(ogen) suggest that t-PA adheres to the fibrin surface with subsequent addition of plasminogen to form a cyclic ternary complex, thus preventing the activation of circulating plasminogen (151). The plasmin formed on the fibrin surface is not readily inactivated by circulating  $\alpha_2$ -antiplasmin (half-life 10-100 seconds). Circulating plasmin, on the other hand, has a half-life of only 0.1 seconds (152).

#### *Urokinase-type plasminogen activator (u-PA)*

u-PA is synthesized as a single chain (scu-PA) pro-urokinase, which is converted to the active two-chained urokinase-type plasminogen activator (tcu-PA) by plasmin. Even enzymes such as kallikrein, trypsin, cathepsin B, human T cell associated serine proteinase-1, and thermolysin may catalyze the proteolysis of scu-PA. tcu-PA is a 411 residue serine protease with MW 55 kDa containing the active site triad His<sup>204</sup>, Asp<sup>255</sup>, and Ser<sup>356</sup>. The plasminogen activating potential of scu-PA is small compared with tcu-PA. However, the modest plasmin quantity generated by scu-PA is sufficient to stimulate further plasminogen activation by converting scu-PA to the active tcu-PA (153;154).

Ligation of u-PA to its cell-surface receptor (u-PAR) found in a variety of cells, is important for its activity. Plasmin generated from receptor bound u-PA plays a central role in the mediation of pericellular proteolysis, and is protected from rapid inactivation by its principal circulating inhibitor  $\alpha_2$ -antiplasmin (155).

### ***Inhibition of fibrinolysis***

#### *$\alpha_2$ -antiplasmin*

$\alpha_2$ -antiplasmin is the principal inhibitor of human plasmin (156). The 464 residue serpine is a glycoprotein with MW 67 kDa, and the plasma concentration is approximately 1 mM. The 464 residue  $\alpha_2$ -antiplasmin and a shorter form containing 453 amino acids are

present in plasma in about equal concentrations, and both form stable complexes with plasmin (157). Through interaction with the  $\alpha$ 2-antiplasmin reactive site located at Arg<sup>376</sup>-Met<sup>377</sup> plasmin forms a 1:1 stoichiometric inactive complex (158). Plasmin not neutralized by  $\alpha$ 2-antiplasmin may be inactivated by circulating  $\alpha$ 2-macroglobuline (159).

#### *Plasminogen activator inhibitor*

In addition to inhibition of plasmin, fibrinolysis may also be inhibited at the activator level by plasminogen activator inhibitor 1 (PAI-1). It is a single chain, 379 residue glycoprotein with MW 52 kDa belonging to the serpine family (160;161). The active site is located at Arg346-Met347. PAI-1 becomes activated through interaction with vitronectin. This interaction induces a conformational change and an elongation of the reactive center loop (RCL) of the molecule, thus enabling the active binding site for plasminogen activators (162). Active PAI-1 rapidly interacts with one- and two-chained t-PA and two-chained u-PA, thus efficiently preventing fibrinolysis. In addition to its principal role as inhibitor of activators of fibrinolysis, PAI-1 has been shown to participate in angiogenesis.

#### *Thrombin-Activatable Fibrinolysis Inhibitor*

Thrombin-activatable fibrinolysis inhibitor (TAFI) is a single chain protein in the family of zinc containing metallopeptidases with MW 60 kDa. It is produced in the liver in its inactive form, TAFI, and found in plasma at a concentration of 75 nM. Following activation by thrombin, plasmin or trypsin, the active form, TAFIa, exerts its antifibrinolytic effect by cleaving off C-terminal lysine residues from partially degraded fibrin. The lysine residues stimulate t-PA mediated plasmin activation. TAFIa, thus, prevents additional binding of plasminogen to the clot and inhibits further fibrinolysis (163;164).



**Fibrinogen assays**

Different techniques have been employed to determine the fibrinogen concentration in plasma. The method described by Jacobsson in 1955 has often been regarded as a reference method, reflecting the total clottability of fibrinogen (165). The fibrinogen concentration is measured after adding thrombin to a buffer diluted plasma sample, leaving the sample to clot for 2 hours, with subsequent syneresis and washing of the clot, and measuring the total clottable protein content spectrophotometrically after dissolving the clot in urea. The method is laborious and time consuming, and is not easily automated. Thus, it is not commonly used as a routine method. An assay based on the clotting rate of a diluted plasma sample was described by Clauss in 1957 (166), and implies the addition of a large quantity of thrombin to a diluted plasma sample, and recording the time to clot formation. Due to the high thrombin concentration, fibrin monomers form rapidly, and the method reflects the fibrinogen concentration and the polymerization properties of the fibrin monomers. The coagulation endpoint can be assessed manually as elapsed time to the formation of a visual clot, but clotting rate assays are also easily automated, making them suitable for routine purposes.

Other methods, such as salt or heat precipitation (167-169), are less commonly used as routine assays. Immunochemical assays are based on polyclonal antibodies reacting with epitopes on the fibrinogen molecule (167;170). The light scatter of insoluble antigen-antibody complexes correlates to the fibrinogen concentration in the sample. Other immunological methods include fibrinogen quantification using ELISA techniques. The variety of methods used to determine plasma fibrinogen levels, in addition to heterogeneous calibration standards, have made comparison of fibrinogen levels obtained from different laboratories difficult (171). In order to obtain comparable estimates of fibrinogen concentrations, the First International Fibrinogen Standard (IS) (89/644) was introduced by the National Institute for Biological Standards in 1992 (172), and replaced in 2000 by the 2<sup>nd</sup> IS for fibrinogen (173). The freeze-dried calibration plasma has enabled laboratories and manufacturers to calibrate against a common international fibrinogen standard.

### **Fibrinogen interaction with blood platelets and leukocytes**

Through interaction with specific cell surface integrin receptors, fibrinogen may interact with blood platelets and leukocytes. Integrins are transmembrane, heterodimeric glycoprotein receptors, and their main tasks include coordination of cell adhesion to extra cellular matrix and cell to cell interactions (174). After activation by agonists such as ADP, collagen or thrombin the platelets become activated and display the GP IIb/IIIa integrin ( $\alpha 2b\beta 3$ ), allowing fibrinogen-platelet interaction. Fibrinogen is the predominant ligand for activated platelets at low shear rates, whereas vWF is the physiologically important ligand at high share rates (175;176). The platelet-fibrinogen interaction is closely regulated to prevent spontaneous intravascular aggregation with potential deleterious effects. The IIb/IIIa receptor, therefore, is not constitutively expressed on platelets. Although both the fibrinogen  $\alpha$ -chain Arg-Gly-Asp (RGD)-sequences (A $\alpha$  95-97 and 572-574) and the C terminal  $\gamma$ -chain ( $\gamma$  400-411) are potential ligands for GP IIb/IIIa, fibrinogen-dependent platelet aggregation requires an intact C-terminal  $\gamma$ -chain. Thus, the alternatively spliced  $\gamma$ -chain,  $\gamma'$ , does not support platelet aggregation (177). The  $\alpha$ -chain RGD sequences are neither necessary nor sufficient for platelet aggregation (178). Fibrinogen may also interact with the leukocyte integrin  $\alpha_M\beta_2$  (CD11b/CD18, Mac-1). The fibrinogen  $\gamma$ - chain region ( $\gamma$  377-395 and  $\gamma$  190-202) functions as ligands for the leukocyte  $\alpha_M\beta_2$  in vivo (179;180). Thus, fibrinogen binds to activated leukocytes, in the presence of divalent cations, with the association of about 150 000 fibrinogen molecules per cell (181). Previous studies have shown that fibrin(ogen) may have different effects on leukocyte function (182-184), and fibrin is involved in monocyte adhesion to endothelium and migration at sites of inflammation (185;186). Furthermore, fibrinogen plays an important role in the inflammatory response to implanted biomaterials (187;188). In addition to platelets and leukocytes, fibrinogen may also interact with fibroblasts, endothelial- and, epithelial cells (189;190).

## 5. AIMS OF THE STUDY

- To investigate the effect of freeze-drying on the clotting properties of fibrinogen, and how the use of a freeze-dried calibration standard might affect the results of a commonly used clotting rate fibrinogen assay.
- To determine whether the fibrinogen concentrations obtained by a clotting rate assay deviated from levels found using a method based on total clottability during acute-phase conditions, and to investigate if a discrepancy between the two methods correlates with changes in the HMW-fibrinogen fraction.
- To determine the influence of the major fibrinogen subfractions on the viscosity of native fibrinogen, and investigate the effect of EDTA on the viscosity of purified fibrinogen and plasma.
- To study the effects of moderate red wine consumption on plasma viscosity, fibrinogen concentration, and the distribution of the 3 major fibrinogen subfractions.
- To investigate potential proinflammatory effects of purified fibrinogen and fibrin on isolated peripheral blood mononuclear cells (PBMC) as reflected by increased mRNA expression and synthesis of the proinflammatory cytokines IL-6, TNF- $\alpha$  and IL-1 $\beta$ , and to assess the significance of an altered HMW/LMW-fibrinogen ratio.

## 6. MATERIALS AND METHODS

### Study subjects

In paper I blood samples drawn from 14 healthy subjects (7 men, 7 women, and median age 22) were used to determine the fibrinogen concentrations in plasma before and after freeze-drying. Blood samples for the preparation of fresh-frozen and freeze-dried calibration standards were collected from 20 healthy subjects (10 men, 10 women, median age 22). The fibrinogen concentration in blood samples from 30 apparently healthy subjects (15 men, 15 women, median age 24) and 10 patients undergoing major elective surgery (4 men, 6 women, median age 66) were used to determine any discrepancy in fibrinogen concentration using the fresh frozen calibration standard compared to the freeze-dried standard.

In paper II the fibrinogen concentrations and the distribution of fibrinogen subfractions in 7 patients about to undergo elective surgery were studied (4 men, 3 women, median age 66 years). Three patients were undergoing total hip joint replacement, 2 had an explorative laparotomy, and 2 patients were undergoing hemicolectomy.

In paper IV we studied the effect of moderate red wine on fibrinogen levels, the distribution of fibrinogen subfractions and plasma viscosity in 92 healthy, nonsmoking men (35%) and women (65%) (mean age 50.3 years). Baseline characteristics of the study population are specified in paper IV.

In paper V we studied the effect of purified fibrinogen and HMW-fibrinogen on the synthesis of proinflammatory cytokines from isolated PBMC collected from 9 healthy subjects (5 women, 4 men, median age 31 years).

### Fibrinogen assays

#### *Total clottable protein assay*

The total clottable protein assay was performed according to a modification of the method described by Jacobsson (1965). Plasma was diluted to one fourth of its original concentration by adding phosphate buffer and thrombin (final concentration 7.5 NIH U/ml). After clotting for 30 min, the clot was synerized, washed in 0.9% saline and dissolved in 40% alkaline urea. The fibrinogen concentration was then calculated by measuring the absorption at 280 nm using the clot solvent as a blank, and correcting for extraneous absorption at 320 nm and 360 nm. The intra-assay coefficients of variation were 1.5% at a fibrinogen concentration of 3.6 g/l (n = 10) and 1.6% at a concentration of 4.7 g/l (n = 10).

#### *Clotting rate assay*

Fibrinogen concentrations in the experiments presented in paper II and III were determined using a clotting rate assay as described by Clauss (1991). 200 µl of a solution containing 1 volume citrated plasma and 9 volumes sodium barbital buffer were incubated at 37 °C for 2 minutes. 100 µl thrombin 100 NIH U/ml was added, and the clotting time was determined. If the clotting time was shorter than 7 seconds, a higher dilution of plasma was used. If the clotting time exceeded 15 seconds, a lower plasma-dilution was used.

#### *Automated clotting rate assay*

Fibrinogen concentrations in the studies presented in paper I and IV were determined by and automated clotting-rate assay (Automated Coagulation Laboratory Futura®, Instrumentation Laboratory, Lexington, MA, USA) using the principle of turbidimetric clot detection. The intra-assay CV for the clotting-rate assay was 3.2% (n=10)

#### ***Freeze-drying***

Freeze-drying of plasma samples (1 ml in each sample) was performed by The Axis-Shield company (Oslo, Norway), based on international guidelines (1992), using a Gambolt GT 150 freeze dryer. The samples were sealed under vacuum, stored in small glass ampoules at -20 °C and reconstituted by adding 1 ml of purified water before use. Residual humidity in freeze-dried plasma samples was 0.4–0.6%.

***Thrombin clotting time***

0.1 ml of 3 IU/ml thrombin was added to 0.2 ml plasma or fibrinogen solution, incubated at 37 °C and the clotting time recorded.

***Electrophoresis***

Gels (3% polyacrylamid and 0.5% agarose) were prepared as described by Weinstein and Deykin (193). The electrophoretic procedure was performed using a Mini-PROTEAN® II (BioRad, Richmond, CA, USA). The gels were Coomassie-stained, and the fibrinogen fractions quantified by densitometric scanning using Kodak image station 440 CF with Kodak 1D image analysis software. The intra-assay coefficient of variation for the electrophoresis and densitometric scanning was 1.8% (n=10).

***Fibrinogen purification***

Fibrinogen used in paper III was purified from plasma donated by healthy volunteers using the  $\beta$ -alanine precipitation method described by Jacobsen and Kierulf (194), except that citrate (final concentration 0.011 M) was present during the entire procedure. Trasylol (final concentration 200 KIU/ml) and soybean trypsin inhibitor (STI, final concentration 0.1 mg/ml) were used as inhibitors instead of Epsilon aminocaproic acid.

***Separation of HMW-, LMW- and LMW'-fibrinogen***

Separation of the fibrinogen fractions was performed by gradual precipitation of a purified fibrinogen solution. A saturated ammonium sulphate solution was added slowly during constant stirring. The HMW-fraction was prepared by the addition of saturated ammonium sulphate to 19% saturation. After centrifugation at 1200 g for 7 min the precipitate was collected, dissolved in 0.15 M NaCl at 37 °C, re-precipitated in ammonium sulphate (19%), leaving the precipitate to dissolve in 1/3 of its original volume and finally dialyzed. The LMW-fraction was obtained from precipitate formed at 22–24% ammonium sulphate saturation and the LMW'-fraction at 26–30% saturation. Precipitates were centrifuged, dissolved in 0.15 M NaCl at 37 °C and dialyzed. All fractions were dialyzed three times against 500 vol diemal buffered saline for 24 h.

Contamination with soluble fibrin and other low solubility complexes were removed by leaving the dialyzed fractions at + 4 °C for 24 hours and removing insoluble material after centrifugation. The purity of the fibrinogen fractions was determined by sodium dodecyl sulphate (SDS) gel electrophoresis. The HMW-fibrinogen fraction was contaminated with 1% LMW. The LMW-fraction contained 96% LMW- and 4% LMW'-fibrinogen, and the LMW'-fraction contained 90% LMW'-fibrinogen and 10% HMW- and LMW-fibrinogen. The thrombin clotting times were 15, 24 and 31 seconds for HMW-, LMW- and LMW'-fibrinogen, respectively. The clottability of HMW-fibrinogen was 100%, of LMW-96% and of LMW'-fibrinogen 76%.

### ***Viscometry***

The viscosity of fresh plasma, purified fibrinogen and fibrinogen subfractions were determined using Haake Microvisco<sup>®</sup> 2 viscometer (Haake Mess-Technik, Munich, Germany). A metal sphere of standard weight and size was inserted into a precision cut syringe containing the sample, and placed in a tilted position inside the viscometer containing two photo sensors. The time needed by the ball to pass between the two photo sensors was measured and the viscosity calculated. The intra-assay coefficient of variation for the viscometer was 0.4% (n=10).

### ***Isolation of PBMC***

Whole blood was drawn by clean venipuncture from healthy volunteers who had not used any medication two weeks prior to inclusion, and collected into lipopolysaccharide (LPS)-free Vacutainer<sup>®</sup> tubes (Cedex, France) containing 7 mM EDTA. PBMC were isolated from whole blood under sterile conditions using Lymphoprep<sup>®</sup> (Axis-Shield PoC, Norton, MA, USA). Cell cultures comprised 81% lymphocytes, (25, 75 percentiles: 69, 82%), 11% monocytes (11, 14%) and 8% granulocytes (7, 16%).

### ***RNA isolation and mRNA quantification***

PBMC were incubated with purified HMW- or unfractionated fibrinogen (final concentration 0.2 mg/ml or 0.5 mg/ml) and RPMI-1640 containing 25 mM HEPES and L-glutamine. Following centrifugation the cell pellets were lysed in 250 µl lysis/binding

buffer (MagNA Pure LC RNA Isolation Kit-High Performance) and stored at  $-70^{\circ}\text{C}$  until analyzed. Total RNA was isolated using a MagNA Pure robot (Roche Diagnostics, Basel, Switzerland) and the “RNA hp-cells” protocol. The amount of total RNA was quantified using the NanoDrop<sup>®</sup> spectrophotometer (NanoDrop technologies, Wilmington, DE, USA) as described in paper V. Detection of the relative change in gene expression in fibrinogen stimulated PBMC compared to controls normalized to an endogenous reference (GAPDH mRNA) was estimated using the Comparative C Method of relative quantification ( $\Delta\Delta\text{Ct}$  method) (195).

### ***Cytokine analysis***

Concentrations of IL-6, TNF- $\alpha$  and IL-1 $\beta$  were determined in cell-lysate using commercial enzyme linked immunosorbent assays (ELISA) from R&D Systems (Minneapolis, MN, USA). Measurements were done in duplicates according to the manufacturer’s instructions and the mean concentrations were calculated. The intra-assay CVs for the IL-6 cytokine assay were 1.7% at low (16.8 pg/ml) and 4.4% at high (191 pg/ml) concentrations, whereas the inter-assay CVs were 2.0% and 3.7%, respectively. Intra-assay CVs for the IL-1 $\beta$  assay were 2.4 at low (12.6 pg/ml) and 3.4% at high (192 pg/ml) concentrations, and the inter-assay CVs 3.4% and 7.1%, respectively. For the TNF- $\alpha$  assay the intra-assay CVs were 5.3% at low (2.4 pg/ml) and 8.8% at high (14 pg/ml) concentrations, and the inter-assay CVs 10.8% (2.4 pg/ml) and 16.7% (14 pg/ml), respectively.

### ***Statistics***

Normally distributed data were reported as mean  $\pm$  standard deviation (SD). Comparison of data sets were valuated using Student’s t-test, and reported as the mean difference with 95% confidence interval (paper IV). Correlations between normally distributed variables were determined using Pearson’s method.

If not normally distributed, data were presented as medians with range or the 25, 75 percentile. Comparisons between data sets were made using Wilcoxon signed rank test (paper I, II, III, V), and correlations between variables performed using Spearman’s rank correlation test. P values are two sided, and values less than 0.05 were regarded as



statistically significant. Statistical analyses were performed using the Statistical Package for Social Sciences version 11.0 and SamplePower 2.0 (SPSS Inc., Chicago, Illinois, USA).

## 7. SUMMARY OF RESULTS

### I

#### **Influence of freeze-drying on the clotting properties of fibrinogen in plasma**

In paper I we studied the effect of freeze-drying on the clotting properties of fibrinogen. Freeze-dried plasma standards are commonly used to calibrate fibrinogen assays. Altered clotting properties following the freeze-drying process might influence the results obtained when using clotting rate assays to determine fibrinogen concentrations. To evaluate the effect of freeze-drying, we determined the fibrinogen concentrations in citrated plasma samples using a total clottable protein method and a clotting rate assay before and after freeze-drying. When using the clotting rate assay, significantly higher fibrinogen concentrations were found in fresh-frozen plasma samples compared to freeze-dried samples ( $P < 0.001$ ). In freeze-dried plasma samples, the fibrinogen concentrations were significantly higher using the total clottable protein assay than the clotting rate assay ( $P < 0.001$ ). When measuring the fibrinogen concentrations in plasma samples with a wide range of fibrinogen concentrations using the clotting rate assay, significantly higher fibrinogen concentrations were found using the freeze-dried calibration plasma, than the fresh-frozen calibration plasma ( $P = 0.02$ ). The results indicate that the clotting rate of fibrinogen in citrated plasma is reduced following freeze-drying. When applying freeze-dried calibration plasma in a clotting rate assay, higher fibrinogen concentrations are obtained than by using fresh-frozen plasma.

### II

#### **Discrepancy between fibrinogen concentrations determined by clotting rate and clottability assays during the acute-phase reaction**

Little is known about the influence of the acute-phase reaction on clotting rate assays. In this paper we wanted to investigate discrepancies between fibrinogen concentrations obtained by a clotting rate assay (as described by Clauss) and a total clottable protein assay (according to Jacobsson). We compared the fibrinogen concentrations using the two methods, and determined the HMW (High Molecular Weight)-, LMW- and LMW'-fibrinogen fractions in plasma samples collected preoperatively and on postoperative days 1, 3, and 5 in patients undergoing major elective surgery. Good agreement between

the two assays was found in samples drawn before the operation. In samples collected on postoperative days 1 and 3, the fibrinogen concentrations determined by the clotting rate assay were significantly higher than the concentrations measured with the total clottable protein assay ( $p = 0.015$  on both days). SDS-gel electrophoresis showed an increase in the median HMW-fraction from 69.7% (range 64.3-70.4) in preoperative samples to 85.8% (80.7-87.6) in samples drawn on day 3. The difference between fibrinogen concentrations obtained by the two methods was significantly correlated to the HMW-fraction of the samples ( $r=0.81$ ,  $p < 0.0001$ ).

### III

#### **The viscosity of fibrinogen subfractions and of EDTA denatured fibrinogen do not differ from that of native fibrinogen**

Fibrinogen is a major determinant of plasma viscosity. It has been suggested that the increased risk of cardiovascular disease associated with a high fibrinogen concentration may partly be attributed to its effect on viscosity. In paper III we wanted to evaluate how the three fibrinogen subfractions high molecular weight (HMW)-, low molecular weight (LMW)-, and very low molecular weight (LMW<sup>-</sup>)-fibrinogen affect viscosity. Furthermore, the viscosity of plasma is usually determined in ethylenediaminetetraacetic acid (EDTA) anticoagulant plasma samples. EDTA affects the clotting properties of fibrinogen due to denaturation, which might also affect the viscosity of fibrinogen. Thus, we also wanted to investigate the effects of EDTA on the viscosity of fibrinogen. Purified fibrinogen was obtained by beta-alanine precipitation of plasma from healthy donors. Separation of the fibrinogen fractions was performed by gradual precipitation of purified fibrinogen by ammonium sulphate. The viscosity was determined using a Haake Microvisco 2 viscometer. The viscosity of the three fibrinogen subfractions did not differ significantly from that of native fibrinogen. A significant prolongation of the thrombin clotting time was observed in the fibrinogen solution containing EDTA at 37 °C compared to 20 °C. However, the viscosity of EDTA anticoagulated purified fibrinogen and plasma samples did not differ from that of heparin anticoagulated samples.

#### IV

##### **A daily glass of red wine induces a prolonged reduction in plasma viscosity: a randomized controlled trial.**

Moderate red wine consumption has been linked to reduced risk of coronary heart disease. Reduced plasma viscosity and reduced fibrinogen levels have been launched as possible contributors to this risk reduction. In study IV we investigated the effect of moderate red wine consumption on plasma viscosity, fibrinogen concentration and fibrinogen subfractions. We recruited 92 healthy, nonsmoking volunteers to consume one glass of red wine daily for 3 weeks in a prospective, randomized cross-over study. In the second 3-week period the volunteers abstained from alcohol use. The plasma viscosity, fibrinogen concentration and the distribution of the main fibrinogen subfractions were determined at inclusion, after wine drinking and after abstinence. Plasma viscosity was reduced by 0.026 and 0.024 mPa.s in the two groups following wine intake (95% CI 0.009-0.043,  $p=0.004/0.0083-0.039$ ,  $p=0.003$ ). The decrease in plasma viscosity was maintained following 3 weeks of abstinence. The fibrinogen concentration was reduced by 0.17 g/l following wine drinking in the group starting with abstinence (CI 0.04-0.29,  $p=0.01$ ). The distribution of the fibrinogen subfractions remained unaltered. The decreased viscosity levels were maintained after 3 weeks of abstinence, suggesting a sustained viscosity lowering effect of red wine.

#### V

##### **Fibrinogen and fibrin induce synthesis of proinflammatory cytokines from isolated peripheral blood mononuclear cells**

Fibrinogen in plasma includes three main fractions; high-molecular-weight (HMW) -fibrinogen, low-molecular-weight (LMW) -fibrinogen, and very-low-molecular-weight (LMW') -fibrinogen. During acute-phase conditions, plasma fibrinogen levels and the HMW-/LMW-fibrinogen ratio increase rapidly due to increased synthesis of HMW fibrinogen. The consequences of elevated plasma fibrinogen levels and local deposition of fibrin in inflammatory tissues observed during acute-phase conditions are not clear. We wanted to investigate proinflammatory effects of fibrinogen and fibrin on peripheral blood mononuclear cells (PBMC) as reflected by altered mRNA expression and synthesis

of the proinflammatory cytokines IL-6, TNF-  $\alpha$  and IL-1 $\beta$  , and to explore the significance of altered HMW-/LMW-fibrinogen ratio. PBMC were isolated from whole blood using Lymphoprep<sup>®</sup>. HMW-fibrinogen was separated from unfractionated fibrinogen by ammonium sulphate precipitation. Cells were incubated with unfractionated fibrinogen, HMW-fibrinogen or fibrin. Cytokine levels in cell lysates were determined using ELISA assays. Real-time PCR was used for mRNA quantification. We found that fibrinogen significantly increased mRNA levels, and induced synthesis of the proinflammatory cytokines IL-6 and TNF-  $\alpha$  in PBMC in a dose dependent manner. Median (25, 75 percentile) IL-6 and TNF-  $\alpha$  concentrations were 12 (5, 40) pg/ ml and 16 (0,61) pg/ml, respectively. Median mRNA quantity was increased 12.3– (6.6, 48.6) and 1.7– (1.5, 6.5) fold for IL-6 and TNF-  $\alpha$  compared to controls. The stimulatory effect of unfractionated fibrinogen was not significantly different from HMW-fibrinogen. Fibrinogen and fibrin were equally effective in promoting cytokine synthesis from PBMC.

## 8. GENERAL DISCUSSION

### Paper I

In 1991 the standardization and scientific committee of the International Society on Thrombosis and Haemostasis (ISTH) supported the establishment of a lyophilized fibrinogen standard (coded 89/644) as an international standard for fibrinogen. The lyophilized plasma was manufactured by The National Institute for Biological Standards and Control (UK) and comprised lyophilized material from a plasma pool of 25 healthy donors (196). The fibrinogen concentration was determined by 22 different laboratories, and the declared concentration of 2.4 g/l clottable protein was the geometric mean of the reported values from the participating laboratories. A total clottable protein method was the recommended assay to determine the fibrinogen concentration in the standard. The use of a common calibration standard is essential to obtain accurate and comparable results between laboratories. Previous studies have indicated that the declared value of standards in commercial available fibrinogen kits may vary by as much as 200% (197). In paper I we investigated the effect of freeze-drying of plasma samples on the fibrinogen concentrations obtained by an modified automated Clauss assay compared to a total

clottable protein according to Jacobsson (165). The total clottable protein assay yielded similar results in fresh frozen and freeze-dried samples. When using the clotting-rate assay, however, significantly higher fibrinogen concentrations were found in fresh-frozen samples compared to those found in freeze-dried specimens (mean difference 0.18 g/l, 95% confidence interval 0.13– 0.23,  $P < 0.001$ ). The precision of both fibrinogen assays were evaluated, and given as the percent coefficient of variation (CV). The intra-assay CVs of the fibrinogen assays used were less than 2 % at the fibrinogen concentrations found in our samples, which indicate a high precision. Although inter-assay CVs were not determined, between-days variability was reduced by performing all comparisons between the two assays on the same day and by the same operator. Freeze-drying has been suggested to induce changes in the secondary and tertiary structure of proteins (198;199). This denaturation might affect its physical properties and biological activity. As the principal function of fibrinogen is clotting, the freeze drying process may inflict conformational changes in the native structure which might affect its clotting properties. Reduced clottability has been considered as evidence of denaturation of fibrinogen. Ly and coworkers suggested, however, that the thrombin clotting time might be a better indicator of denaturation than the clottability (200). They also found that the denaturation process did not reduce the fibrinogen N-terminal susceptibility to thrombin. This may imply that denaturation primarily affects the polymerization properties of the fibrin monomers, thus, influencing assays based on clotting rate to a larger extent than assays reflecting the total clottability of fibrinogen. The use of diluted plasma and large amounts of thrombin in the clotting rate method used in our study makes this assay even more sensitive to polymerization defects than the thrombin clotting time. The freeze-drying process may also alter the turbidity of the plasma samples (173), which may induce discrepancies in results obtained by the two assays. The fibrinogen concentration is calculated after spectrophotometrically measuring the absorption at 280 nm and correcting for extraneous absorption at 320 nm and 360 nm. Samples with optical densities (OD) at 320 nm exceeding 10% of OD 280 may underestimate the fibrinogen concentration. Freeze drying of the plasma samples had little effect of the turbidity in our study ( $OD_{320} < 10\%$  of  $OD_{280}$ ), and the freeze drying process did not affect the results obtained by the clottability assay compared to fresh frozen samples. To investigate the

effect of using a freeze-dried plasma standard compared to a fresh-frozen standard, the fibrinogen concentrations in plasma samples from individuals with a wide range of fibrinogen concentrations were determined using the clotting-rate assay. Significantly higher fibrinogen concentrations were obtained using the freeze-dried calibration plasma compared to corresponding fresh-frozen plasma (mean 0.04 g/l, 95% CI 0.006-0.08,  $p=0.02$ ). The difference is small, however, and probably negligible in every-day clinical practice. In epidemiological studies, however, when comparing minor differences in fibrinogen concentrations, the influence of using freeze-dried calibration plasmas should be taken into consideration. The 1<sup>st</sup> fibrinogen standard was replaced in 2000 by the 2<sup>nd</sup> International Standard for Fibrinogen (coded 98/612) prepared by the UK NIBSC and recommended by the ISTH fibrinogen subcommittee. The fibrinogen content of this standard was determined using an automated clotting-rate assay (173).

## **Paper II**

In paper II we compared the plasma fibrinogen concentrations in patients undergoing major elective surgery obtained by a commonly used clotting rate assay (166), with a total clottable protein assay according to a modification of the method described by Jacobsson (165). Holm and Godal had previously shown that the HMW-/LMW-fibrinogen ratio increases rapidly during acute-phase conditions such as myocardial infarction and major surgery (201), and that the thrombin clotting times of LMW- and LMW'-fibrinogen were prolonged compared to HMW-fibrinogen due to altered polymerization properties of the 3 fibrinogen fractions (63). The clotting rate assay procedure implies the addition of a high concentration of thrombin (final concentration 33 NIH U/ml), with a subsequent rapid release of fibrinopeptide A (FPA), resulting in a clotting rate depending on fibrinogen concentration and the polymerization properties of the fibrin monomers. A high correlation between the divergence in fibrinogen concentrations obtained by the two methods and the HMW-fibrinogen level was found ( $r = 0.81$ ), indicating that the increase in HMW-fibrinogen may be an important contributor to the observed discrepancy.

Both fibrinogen assays used in the study have been extensively validated previously (165;166), and the precision of the two assay was good (intra assay CVs were 1.2-3.2% and 1.5-1.6% for the clotting rate and the clottability assays respectively). All samples

from each patient were assayed on the same day and by the same operator, thus reducing between-days variability. Acute-phase conditions affects the phosphorylation of FPA, which may increase from the normal 25-30% to over 50% (141). This has been suggested to affect the release rate of FPA. However, a more effective release of phosphorylated FPA than of FPA would probably have a limited effect on our results since high thrombin concentrations are added to diluted plasma. The blood samples were obtained from non-fasting patients. Although it is clear that lipoproteins are included in fibrin clots obtained from lipemic plasma (165), the error in fibrinogen determination of lipemic plasma samples is small, providing that correction for "extraneous absorption" is made (165). All clottability measurements in our study were done at 280 nm wavelength, correcting for extraneous absorption at 320 and 360 nm. Furthermore, all plasma samples were visually inspected and none found to be lipemic.

The results illustrate that variations in the HMW-/LMW-fibrinogen ratio may affect results obtained by a clotting rate assay, and contribute to discrepancies between methods reflecting the total clottability of fibrinogen and assays reflecting the functional state of fibrinogen.

### **Paper III**

In paper III we investigated the influence of the 3 major fibrinogen subfractions on the viscosity of native fibrinogen, and found that the viscosity of high molecular weight (HMW-) fibrinogen did not differ from that of low molecular weight (LMW)-, or very low molecular weight (LMW') fibrinogen. This implies that degradation of the C-terminal end of one- (LMW-fibrinogen) or both (LMW'-fibrinogen)  $\alpha$ -chains of the fibrinogen molecule, representing a loss in molecular weight of 35 kDa (10% of native weight) and 70 kDa (20 % of native weight,) respectively, does not significantly reduce the viscosity of fibrinogen. Fibrinogen is a major determinant of plasma viscosity (134;135). Thus, altered plasma viscosity attributable to increased fibrinogen levels in acute-phase conditions is not related to concomitant changes in the distribution of the major fibrinogen subfractions. At equal concentrations the size and shape of proteins determine their effect on plasma viscosity. Normally, the  $\alpha$ C domain of the fibrinogen molecule extends from the D region, parallel to the coiled-coil of the A $\alpha$ -, B $\beta$ - and  $\gamma$ -



chains, interacting with the central region of the molecule or forming intra-molecular  $\alpha$ C- $\alpha$ C connections (29;30). Following cleavage of FPB the  $\alpha$ C domains dissociate from the central region and become available for inter-molecular  $\alpha$ C- $\alpha$ C connections (30). The lack of effect on fibrinogen viscosity following the loss of the  $\alpha$ C-domains may indicate that the  $\alpha$ C-domain is of minor importance to the overall shape of the fibrinogen molecule. Potential sources of error in this study comprise impurities of the isolated fibrinogen fractions. Separation of the fibrinogen fractions was performed by gradual precipitation of a purified fibrinogen solution, and the purity of the subfractions determined by sodium dodecyl sulphate (SDS) gel electrophoresis. The HMW-fibrinogen fraction was contaminated with 1% LMW. The LMW fraction contained 96% LMW- and 4% LMW'-fibrinogen, and the LMW' -fraction contained 90% LMW'-, and 10% HMW- and LMW-fibrinogen. The reason why the LMW'-fibrinogen contained HMW-fibrinogen and the LMW-fibrinogen fraction did not is unclear. Possibly, the higher ammonium sulphate concentration needed to precipitate the LMW'-fibrinogen fraction may have promoted a more effective precipitation of the remaining HMW-fibrinogen in the solution. Contamination of the fibrinogen solutions with soluble fibrin and other low solubility complexes were reduced by leaving the dialyzed fractions at 4 °C for 24 h and removing insoluble material following centrifugation. Remaining contaminations with proteins such as fibronectin represent potential sources of error. Fibronectin is a common contaminant in purified fibrinogen solutions, and it is up-concentrated during the fibrinogen precipitation procedure (202). The fibronectin concentrations in the fibrinogen subfractions in our study were low; 1.8  $\mu$ g/g fibrinogen (HMW fraction), 6.0  $\mu$ g/g fibrinogen (LMW fraction) and 26  $\mu$ g/g fibrinogen (LMW' fraction). Furthermore, the specific viscosity of fibrinogen is significantly higher than that of other plasma proteins. Thus, protein contamination of the LMW'-fibrinogen fraction would reduce the viscosity. No differences in viscosity levels in the 3 fibrinogen fractions were observed in our study. The intra-assay CV for the quantification of the fibrinogen fractions was 1.8%, indicating a high precision for the electrophoresis and densitometric scanning procedures. Adding EDTA, a commonly used anticoagulant, to purified native fibrinogen did not affect the viscosity despite a significant increase in thrombin clotting time. EDTA is a chelating agent which effectively removes calcium. Calcium ions are essential to preserve

the structure and function of the fibrinogen molecule (200;203;204). Removal of calcium facilitates denaturation of fibrinogen as reflected by increased thrombin clotting time (33;200). Denaturation of proteins has been closely associated with changes in viscosity (205), but the denaturation of fibrinogen induced by EDTA does not significantly affect the viscosity of plasma compared to heparin anticoagulated plasma.

#### **Paper IV**

In paper IV we wanted to investigate the effect of a daily glass of red wine on plasma viscosity, fibrinogen levels and the distribution of the major fibrinogen subfractions. Moderate alcohol consumption has been linked to reduced risk of ischemic heart disease (206-208). A beneficial effect on haemostatic factors such as fibrinogen and plasma viscosity has been launched as possible contributors to this risk reduction. The study was a prospective, randomized crossover trial, and a sub-study of the work by Retterstøl and colleagues from 2005 (209). The study recruited 92 healthy men (35%) and women (65%). Smoking significantly affects fibrinogen levels and plasma viscosity, and only non-smokers were recruited in the trial (210). The crossover design implies that the volunteers were randomized to start with 3 weeks of red wine drinking or abstinence, and subsequently crossed over to the other intervention. This design is particularly effective in intervention studies, because it enables each individual to be its own control, and, thus, eliminates between-subject variation. This increases the statistical power of the study compared to parallel group designs, and requires a smaller sample size. Crossover studies are, however, sensitive to carryover effects. As illustrated in figure 1 and 2 in paper IV, the observed mean plasma viscosity and fibrinogen concentration following the abstinence period differed in the subgroups, indicating a prolonged effect of wine drinking on plasma viscosity. This effect and the fact that the majority of the participants were moderate alcohol consumers at the time of inclusion (median consumption 1-3 units per week) may explain why no difference in plasma fibrinogen levels were observed in the group starting with wine drinking. The prolonged effect of wine drinking on plasma viscosity was an unexpected but interesting finding in the study, which, to our knowledge, has not previously been discussed. Including abstinence periods at the beginning of the study and between each treatment period, might have reduced this

effect. Due to the prolonged viscosity lowering effect of red wine, the results from each subgroup were analyzed and presented separately. For logistical reasons, the alcohol intake in the study was not supervised. Although the importance of complying with the study protocol was carefully explained to the participants, lack of adherence to the scheduled intervention represents a potential confounder in the study. Self reported compliance was, however, good. Five participants reported missing 1-2 days in the wine period, and two participants consumed 1-2 units of alcohol during the abstention period, which may have attenuated the effects on viscosity and fibrinogen levels, and potentially reduced the observed differences between the wine drinking- and abstention periods.

Although the participants were encouraged to maintain a normal diet, data concerning diet and physical exercise were not recorded. However, only ethanol consumption seems to have a clear and consistent effect on plasma fibrinogen levels. Other dietary components seem to have modest or no effects (for review see (211)). Thus, changes in diet during the study period do probably not represent a potential confounder on fibrinogen levels. Blood sampling were performed in non-fasting subjects. The effects of meals seem to have little effect on plasma viscosity measurements, however (212;213).

Plasma viscosity was determined by a “falling ball” viscometer. The intra assay CV of the assay was 0.41%, indicating a high precision in measurements. Blood sampling and measurements were performed according to international guidelines, except that frozen plasma samples were used. No statistically significant differences in viscosity between corresponding fresh and thawed frozen plasma samples were found in our study. Differences in study populations, smoking and alcohol consumption, fibrinogen levels and different assays used for viscosity determination, may contribute to differences in observed plasma viscosity in different trials (214-216). Increased plasma viscosity and fibrinogen levels have been identified as independent markers of increased risk of CHD (217). In addition to raising plasma high density lipoprotein (HDL) levels, reduced plasma viscosity and fibrinogen levels may contribute to the observed health benefits associated with moderate red wine consumption.

## Paper V

Close interaction exists between the inflammatory- and coagulations systems. Thus, local and systemic inflammation may induce initiation of the coagulation system, and activation of coagulation may, on the other hand, affect inflammatory activity (for review see (218)). Tissue factor (TF) plays a central role in the inflammation induced activation of coagulation, but the interactions between the two systems also include thrombin, protein C and the fibrinolytic system. On the other hand, proinflammatory cytokines and chemokines have been shown to promote the expression of TF and induce coagulation (219).

In paper V we found that purified fibrinogen and fibrin significantly upregulated mRNA and synthesis of the proinflammatory cytokines IL-6 and TNF- $\alpha$  in isolated peripheral blood mononuclear cells (PBMC). Fibrinogen and fibrin were equally effective, and no difference in stimulatory effects were found between unfractionated fibrinogen containing 59% HMW-, 39% LMW- and 2% LMW'-fibrinogen and purified HMW-fibrinogen containing 99% HMW- and 1% LMW-fibrinogen, suggesting that the C-terminal  $\alpha$ -chain is not essential for the stimulatory effects. Altered HMW-/LMW ratios, thus, seem to be of minor importance for the potential proinflammatory effects of fibrinogen/fibrin. Higher fibrin(ogen) concentrations in the cell cultures yielded higher cytokine levels in the supernatant, indicating a dose dependant effect.

Previous reports have indicated that fibrinogen may affect leukocyte function (182;183;220-224), and Perez and coworkers found that fibrin and fibrinogen may induce synthesis of IL-1 $\beta$  from PBMC (184). The integrins CD11b/CD18 (Mac-1) and, to a less extent, CD11c/CD18, seem to be the principal fibrinogen receptors on monocytes, macrophages and lymphocytes. The fibrin(ogen)  $\gamma$ -chain ( $\gamma^{190-202}$  and  $\gamma^{377-395}$ ) contains the putative recognition sites for the integrins (225). Fibrinogen mediated activation of transcriptional factors such as NF-kappa B have been implicated in the activation process (226;227). Circulating PBMC are continuously exposed to fibrinogen in vivo without producing significant levels of proinflammatory cytokines in healthy individuals (228). Thus, cell activation seems to be required for the cells to express CD11b/CD18 and thus facilitate cell-fibrinogen interaction (228;229).

We used cultured PBMC isolated from whole blood to study the effects of fibrinogen and fibrin. PBMC includes monocytes and lymphocytes and are commonly used to study the effect of stimulants *ex vivo*. The advantage of using isolated cell cultures is that the effects of stimulatory or inhibitory agents on specific cells can be studied in detail. Since cytokines are mainly released and consumed locally at the site of inflammation, cytokine levels in peripheral blood often do not reflect local inflammation and, thus, can not be used to study local inflammatory mechanisms. The disadvantages of using isolated cell cultures include removal of cytokines and other stimulatory or inhibitory factors during the isolation process. These factors may play essential roles in the inflammatory response *in vivo*. Furthermore, the isolation process and subsequent incubation may in it self activate the cells and make them more susceptible to stimulants (230). Therefore, inflammatory responses to stimulants observed in isolated cell cultures do not necessarily reflect the physiological responses *in vivo*. Whole blood systems may be better in reproducing the natural cell environment, allowing interaction between different cells, and including natural occurring stimulatory or inhibitory factors. Several potential confounding factors may have affected the results from our study. Cell activation and synthesis of proinflammatory cytokine due to lipopolysaccharids (LPS) or bacteria contamination of collection tubes, reagents and buffers during the blood sampling or cell isolation process represent an important potential source of error when studying inflammatory responses in cell cultures (231-233). Blood samples in our study were drawn from apparently healthy subjects who had not used any medication two weeks prior to blood sampling. We used sterile, endotoxin tested sampling tubes and buffers, and the PBMC isolation procedure was performed under strict sterile conditions. When present, LPS contamination was removed using an endotoxin removal resin (End-X<sup>®</sup> B-52, Associates of Cape Cod, Woods Hole, MA, USA). Before used, all buffers, culture media, electrolyte solutions and fibrinogen solutions were tested for LPS contamination using a highly sensitive chromogenic end-point assay with a lower detection limit of 1 pg/ml LPS (0.01 EU/ml), and deemed LPS free. LPS contamination below the detection limit of the assay, although not capable of inducing significant cytokine synthesis, may have contributed to cell activation. Following isolation, the cell cultures were incubated at 37 °C for 10 hours, before frozen at -70 °C. Activation of cells during the isolation

procedures and incubation period is well known. Previous trials have demonstrated changes in steady-state expression levels of cytokine mRNAs such as IL-1, IL-6, and TNF- $\alpha$  during culture of human PBMC, even under control conditions (230). Spontaneous in vitro aggregation or adherence to the culture wells might also promote activation and be of relevance in gene expression studies. Cytokine synthesis in our control cell cultures were either absent or weakly detectable. Importantly, however, all cells were exposed to the same isolating and incubation procedures, and supernatant from cells incubated with fibrinogen or fibrin contained significantly higher cytokine levels than controls. Furthermore, incubating cells with liquor from the defibrinated fibrinogen solutions (after clotting with thrombin) did not induce cell cytokine synthesis, indicating that fibrinogen indeed promoted mRNA upregulation and cytokine synthesis from the cells.

Variations in mRNA upregulation and cytokine production in cell cultures from different donors might be attributable to variations in activation or the number of monocytes in the cell cultures. All cell cultures contained significant numbers of granulocytes (25, 75 percentile: 7, 16%), which might also have affected our results. However, previous cell stimulation studies indicate a strong correlation between the production levels of proinflammatory cytokines like IL-1 $\beta$ , IL-6, and TNF- $\alpha$  and the number of monocytes, indicating that monocytes may be the principal source of synthesis of these cytokines in cell cultures (228).

Limited data exists on proinflammatory effects of fibrinogen and fibrin in vivo. It is clear that fibrinogen plays a central role in the inflammatory response to implanted biomaterials (187;188), and animal studies have suggested that local deposition of fibrin sustains inflammation in some inflammatory diseases (234). Due to the close interaction between fibrin(ogen) and inflammatory cells in acute and chronic inflammatory diseases in vivo (235-237), possible proinflammatory effects of fibrin(ogen) might modify the inflammatory responses in a wide range of inflammatory disorders. Further research is needed, however, to elucidate the interaction between fibrin(ogen) and inflammatory cells, and determine potential proinflammatory effects. Experiments should include whole blood systems as well as isolated cell systems including monocytes/macrophages and lymphocyte subclasses.

## 9. CONCLUSIONS

1. Freeze-drying of human plasma decreases its clotting rate. When using a clotting rate fibrinogen assay calibrated with freeze-dried plasma, higher fibrinogen concentrations are obtained compared to using fresh-frozen plasma (paper I).
2. Higher plasma fibrinogen concentrations are obtained using a clotting rate fibrinogen assay than a total clottable protein assay following major surgery. The discrepancy in results from the two methods is correlated to an increased HMW-fibrinogen fraction (paper II).
3. The viscosity of the 3 main fibrinogen subfractions HMW-, LMW-, and LMW'-fibrinogen do not differ from that of native fibrinogen. Adding EDTA to purified fibrinogen induces denaturation of the fibrinogen molecule as reflected by prolonged thrombin clotting time, but does not affect viscosity. The use of EDTA as plasma anticoagulant does not significantly affect plasma viscosity compared to Heparin anticoagulated plasma (paper III).
4. A daily glass of red wine induces a prolonged reduction in plasma viscosity. Fibrinogen levels are also reduced when preceded by an abstention period (paper IV).
5. Fibrinogen upregulates mRNA and synthesis of the proinflammatory cytokines IL-6 and TNF- $\alpha$  in isolated peripheral blood mononuclear cells. Fibrinogen and fibrin are equally effective in promoting cytokine synthesis, and altered HMW-LMW-fibrinogen ratio does not alter the proinflammatory effects (paper V).

## 10. REFERENCES

- (1) Sun NC, McAfee WM, Hum GJ, Weiner JM. Hemostatic abnormalities in malignancy, a prospective study of one hundred eight patients. Part I. Coagulation studies. *Am J Clin Pathol* 1979; 71(1):10-16.
- (2) Holm B, Godal HC. Quantitation of the three normally-occurring plasma fibrinogens in health and during so-called "acute phase" by SDS electrophoresis of fibrin obtained from EDTA-plasma. *Thromb Res* 1984; 35(3):279-290.
- (3) Glenn EM. Fibrinogen and experimental inflammation. *Biochem Pharmacol* 1969; 18(2):317-326.
- (4) de Maat MP. Effects of diet, drugs, and genes on plasma fibrinogen levels. *Ann N Y Acad Sci* 2001; 936:509-521.
- (5) Krobot K, Hense HW, Cremer P, Eberle E, Keil U. Determinants of plasma fibrinogen: relation to body weight, waist-to-hip ratio, smoking, alcohol, age, and sex. Results from the second MONICA Augsburg survey 1989-1990. *Arterioscler Thromb* 1992; 12(7):780-788.
- (6) Meade TW, Imeson J, Stirling Y. Effects of changes in smoking and other characteristics on clotting factors and the risk of ischaemic heart disease. *Lancet* 1987; 2(8566):986-988.
- (7) Margaglione M, Cappucci G, Colaizzo D, Pirro L, Vecchione G, Grandone E et al. Fibrinogen plasma levels in an apparently healthy general population--relation to environmental and genetic determinants. *Thromb Haemost* 1998; 80(5):805-810.
- (8) Sinha S, Luben RN, Welch A, Bingham S, Wareham NJ, Day NE et al. Fibrinogen and cigarette smoking in men and women in the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population. *Eur J Cardiovasc Prev Rehabil* 2005; 12(2):144-150.
- (9) Lee AJ, Lowe GD, Smith WC, Tunstall-Pedoe H. Plasma fibrinogen in women: relationships with oral contraception, the menopause and hormone replacement therapy. *Br J Haematol* 1993; 83(4):616-621.
- (10) Balleisen L, Bailey J, Epping PH, Schulte H, van de LJ. Epidemiological study on factor VII, factor VIII and fibrinogen in an industrial population: I. Baseline data on the relation to age, gender, body-weight, smoking, alcohol, pill-using, and menopause. *Thromb Haemost* 1985; 54(2):475-479.



- (11) Koenig W. Fibrin(ogen) in cardiovascular disease: an update. [Review] [84 refs]. *Thrombosis & Haemostasis* 1996; 76(4):601-9, 2003.
- (12) Chan P, Yeh CJ, Pan WH. Influence of Contraceptive Pill Use, Premenopause and Postmenopausal State on Hemostatic Parameters in Ethnic Chinese Women. *J Thromb Thrombolysis* 1996; 3(1):9-12.
- (13) Meade TW, Haines AP, North WR, Chakrabarti R, Howarth DJ, Stirling Y. Haemostatic, lipid, and blood-pressure profiles of women on oral contraceptives containing 50 microgram or 30 microgram oestrogen. *Lancet* 1977; 2(8045):948-951.
- (14) Rosenson RS, Tangney CC, Mosca LJ. Hormone replacement therapy improves cardiovascular risk by lowering plasma viscosity in postmenopausal women. *Arterioscler Thromb Vasc Biol* 1998; 18:1902-1905.
- (15) Hamsten A, Iselius L, de Faire U, Blomback M. Genetic and cultural inheritance of plasma fibrinogen concentration. *Lancet* 1987; 2(8566):988-991.
- (16) Hall CE, Slayter HS. The fibrinogen molecule: its size, shape, and mode of polymerization. *J Biophys Biochem Cytol* 1959; 5(1):11-16.
- (17) Cohen C. Invited discussion at 1960 symposium of protein structure. *J Polym Sci* 1961; 49:144-145.
- (18) Fowler WE, Erickson HP. Trinodular structure of fibrinogen. Confirmation by both shadowing and negative stain electron microscopy. *J Mol Biol* 1979; 134(2):241-249.
- (19) Weisel JW, Stauffacher CV, Bullitt E, Cohen C. A model for fibrinogen: domains and sequence. *Science* 1985; 230(4732):1388-1391.
- (20) Marchant RE, Barb MD, Shainoff JR, Eppell SJ, Wilson DL, Siedlecki CA. Three dimensional structure of human fibrinogen under aqueous conditions visualized by atomic force microscopy. *Thromb Haemost* 1997; 77(6):1048-1051.
- (21) Caspary EA, Kekwick RA. Some physicochemical properties of human fibrinogen. *Biochem J* 1957; 67(1):41-48.
- (22) Weisel JW. Fibrinogen and fibrin. *Adv Protein Chem* 2005; 70:247-299.
- (23) Henschen A, Lottspeich F. Fibrinogen structure and fibrinolysis. *Haematologica* 1980; 65(4):535-541.
- (24) Doolittle RF. Fibrinogen and fibrin. *Annu Rev Biochem* 1984; 53:195-229.

- (25) Blombäck B, Blombäck M. The molecular structure of fibrinogen. *Ann N Y Acad Sci* 1972; 202:77-97.
- (26) McDonagh J, Messel H, McDonagh RP, Jr., Murano G, Blomback B. Molecular weight analysis of fibrinogen and fibrin chains by an improved sodium dodecyl sulfate gel electrophoresis method. *Biochim Biophys Acta* 1972; 257(1):135-142.
- (27) Blomback B, Blomback M, Henschen A, Hessel B, Iwanaga S, Woods KR. N-terminal disulphide knot of human fibrinogen. *Nature* 1968; 218(5137):130-134.
- (28) Doolittle RF, Goldbaum DM, Doolittle LR. Designation of sequences involved in the "coiled-coil" interdomainal connections in fibrinogen: constructions of an atomic scale model. *J Mol Biol* 1978; 120(2):311-325.
- (29) Gorkun OV, Veklich YI, Medved LV, Henschen AH, Weisel JW. Role of the alpha C domains of fibrin in clot formation. *Biochemistry* 1994; 33(22):6986-6997.
- (30) Veklich YI, Gorkun OV, Medved LV, Nieuwenhuizen W, Weisel JW. Carboxyl-terminal portions of the alpha chains of fibrinogen and fibrin. Localization by electron microscopy and the effects of isolated alpha C fragments on polymerization. *J Biol Chem* 1993; 268(18):13577-13585.
- (31) Hantgan RR, Simpson-Haidaris PJ, Francis CW, Marder VJ. Fibrinogen structure and physiology. In: Colman RW, Hirsh VJ, Marder VJ, Clowes AW, George JN, editors. *Hemostasis and Thrombosis: Basic Principles and Clinical Practice*. Lippincott, Williams and Wilkins, Philadelphia., 2000: 202-232.
- (32) Marguerie G, Chagniel G, Suscillon M. The binding of calcium to bovine fibrinogen. *Biochim Biophys Acta* 1977; 490(1):94-103.
- (33) Ly B, Godal HC. Denaturation of fibrinogen, the protective effect of calcium. *Haemostasis* 1972; 1:204-209.
- (34) Odrlićin TM, Rybarczyk BJ, Francis CW, Lawrence SO, Hamaguchi M, Simpson-Haidaris PJ. Calcium modulates plasmin cleavage of the fibrinogen D fragment gamma chain N-terminus: mapping of monoclonal antibody J88B to a plasmin sensitive domain of the gamma chain. *Biochim Biophys Acta* 1996; 1298(1):69-77.
- (35) Henschen A. Number and reactivity of disulfide bonds in fibrinogen and fibrin. *Arkiv Kemi* 1964; 22:355.
- (36) Townsend RR, Hilliker E, Li YT, Laine RA, Bell WR, Lee YC. Carbohydrate structure of human fibrinogen. Use of 300-MHz <sup>1</sup>H-NMR to characterize glycosidase-treated glycopeptides. *J Biol Chem* 1982; 257(16):9704-9710.

- (37) Langer BG, Weisel JW, Dinauer PA, Nagaswami C, Bell WR. Deglycosylation of fibrinogen accelerates polymerization and increases lateral aggregation of fibrin fibers. *J Biol Chem* 1988; 263(29):15056-15063.
- (38) Drury D, McMaster P. The liver as the source of fibrinogen. *J Exp Med* 1929; 50:569-678.
- (39) Miller LL, Bale WF. Synthesis of all plasma protein fractions except gamma globulins by the liver; the use of zone electrophoresis and lysine-epsilon-C14 to define the plasma proteins synthesized by the isolated perfused liver. *J Exp Med* 1954; 99(2):125-132.
- (40) Collen D, Tytgat GN, Claeys H, Piessens R. Metabolism and distribution of fibrinogen. I. Fibrinogen turnover in physiological conditions in humans. *Br J Haematol* 1972; 22(6):681-700.
- (41) Hart H. The biological half-life of I-131 fibrinogen. *Thromb Diath Haemorrh* 1965; Suppl 27:337-348.
- (42) Straub PW. Chronic intravascular coagulation. Clinical spectrum and diagnostic criteria, with special emphasis on metabolism, distribution and localization of I 131 -fibrinogen. *Acta Med Scand Suppl* 1971; 526:1-95.
- (43) Takeda Y. Studies of the metabolism and distribution of fibrinogen in healthy men with autologous 125-I-labeled fibrinogen. *J Clin Invest* 1966; 45(1):103-111.
- (44) Nham SU, Fuller GM. Effect of fibrinogen degradation products on production of hepatocyte stimulating factor by a macrophage cell line (P388D1). *Thromb Res* 1986; 44(4):467-475.
- (45) Ritchie DG, Levy BA, Adams MA, Fuller GM. Regulation of fibrinogen synthesis by plasmin-derived fragments of fibrinogen and fibrin: an indirect feedback pathway. *Proc Natl Acad Sci U S A* 1982; 79(5):1530-1534.
- (46) Bell WR, Kessler CM, Townsend RR. Stimulation of fibrinogen biosynthesis by fibrinogen fragments D and E. *Br J Haematol* 1983; 54:599-610.
- (47) Franks JJ, Kirsch RE, Frith LO. Effects of fibrinogenolytic products D and E on fibrinogen and albumin synthesis in the rat. *J Clin Invest* 2007; 67:575-580.
- (48) Princer HMG, Moshage H, Emeis JJ. Fibrinogen fragments X, Y, D and E increase levels of plasma fibrinogen and liver mRNA coding for fibrinogen polypeptides in rats. *Thromb Haemost* 1985; 53:212-215.
- (49) Lee SY, Lee KP, Lim JW. Identification and biosynthesis of fibrinogen in human uterine cervix carcinoma cells. *Thromb Haemost* 1996; 75(3):466-470.

- (50) Guadiz G, Sporn LA, Simpson-Haidaris PJ. Thrombin cleavage-independent deposition of fibrinogen in extracellular matrices. *Blood* 1997; 90(7):2644-2653.
- (51) Galanakis D, Nuovo G, Spitzer S, Kaplan C, Scharrer I. Fibrinogen mRNA and antigen co-present in human trophoblasts in situ: possible implications. *Thromb Res* 1996; 81(2):263-269.
- (52) Parrott JA, Whaley PD, Skinner MK. Extrahepatic expression of fibrinogen by granulosa cells: potential role in ovulation. *Endocrinology* 1993; 133(4):1645-1649.
- (53) Kant JA, Fornace AJ, Jr., Saxe D, Simon MI, McBride OW, Crabtree GR. Evolution and organization of the fibrinogen locus on chromosome 4: gene duplication accompanied by transposition and inversion. *Proc Natl Acad Sci U S A* 1985; 82(8):2344-2348.
- (54) Otto JM, Grenett HE, Fuller GM. The coordinated regulation of fibrinogen gene transcription by hepatocyte-stimulating factor and dexamethasone. *J Cell Biol* 1987; 105(3):1067-1072.
- (55) Huber P, Laurent M, Dalmon J. Human beta-fibrinogen gene expression. Upstream sequences involved in its tissue specific expression and its dexamethasone and interleukin 6 stimulation. *J Biol Chem* 1990; 265(10):5695-5701.
- (56) Anderson GM, Shaw AR, Shafer JA. Functional characterization of promoter elements involved in regulation of human B beta-fibrinogen expression. Evidence for binding of novel activator and repressor proteins. *J Biol Chem* 1993; 268(30):22650-22655.
- (57) Zhang Z, Jones S, Hagood JS, Fuentes NL, Fuller GM. STAT3 acts as a co-activator of glucocorticoid receptor signaling. *J Biol Chem* 1997; 272(49):30607-30610.
- (58) Mosesson MW, Finlayson JS, Umfleet RA, Galanakis D. Human fibrinogen heterogeneities. I. Structural and related studies of plasma fibrinogens which are high solubility catabolic intermediates. *J Biol Chem* 1972; 247(16):5210-5219.
- (59) Sherman LA, Fletcher AP, Sherry S. In vivo transformation between fibrinogens of varying ethanol solubilities: a pathway of fibrinogen catabolism. *J Lab Clin Med* 1969; 73(4):574-583.
- (60) Lipinska I, Lipinski B, Gurewich V. Fibrinogen heterogeneity in human plasma. Electrophoretic demonstration and characterization of two major fibrinogen components. *J Lab Clin Med* 1974; 84(4):509-516.

- (61) Gron B, Bennick A, Nieuwenhuizen W, Bjornsen S, Brosstad F. Immunovisualization of fibrinogen A alpha-chain heterogeneity in normal plasma and plasma from patients with DIC or on streptokinase therapy. *Thromb Res* 1988; 52(5):413-424.
- (62) Mosesson MW, Alkjaersig N, Sweet B, Sherry S. Human fibrinogen of relatively high solubility. Comparative biophysical, biochemical, and biological studies with fibrinogen of lower solubility. *Biochemistry* 1967; 6(10):3279-3287.
- (63) Holm B, Brosstad F, Kierulf P, Godal HC. Polymerization properties of two normally circulating fibrinogens, HMW and LMW. Evidence that the COOH-terminal end of the a-chain is of importance for fibrin polymerization. *Thromb Res* 1985; 39(5):595-606.
- (64) Bach-Gansmo ET, Halvorsen S, Godal HC, Skjonsberg OH. Impaired coagulation of fibrinogen due to digestion of the C-terminal end of the A alpha-chain by human neutrophil elastase. *Thromb Res* 1994; 73(1):61-68.
- (65) Phillips HM. Isolation of size homogeneous preparations of high molecular weight and low molecular weight fibrinogens. *Can J Biochem* 1981; 59(5):332-342.
- (66) Sherman LA, Mosesson MW, Sherry S. Isolation and characterization of the clottable low molecular weight fibrinogen derived by limited plasmin hydrolysis of human fraction I-4. *Biochemistry* 1969; 8(4):1515-1523.
- (67) Marder VJ, Shulman NR, Carroll WR. High molecular weight derivatives of human fibrinogen produced by plasmin. I. Physicochemical and immunological characterization. *J Biol Chem* 1969; 244(8):2111-2119.
- (68) Medved LV, Gorkun OV, Manyakov VF, Belitser VA. The role of fibrinogen alpha C-domains in the fibrin assembly process. *FEBS Lett* 1985; 181(1):109-112.
- (69) Medved' LV, Gorkun OV, Manyakov VF, Belitser VA. The role of fibrinogen alpha C-domains in the fibrin assembly process. *FEBS Lett* 1985; 181(1):109-112.
- (70) Fu Y, Weissbach L, Plant PW, Oddoux C, Cao Y, Liang TJ et al. Carboxy-terminal-extended variant of the human fibrinogen alpha subunit: a novel exon conferring marked homology to beta and gamma subunits. *Biochemistry* 1992; 31(48):11968-11972.
- (71) Fu Y, Grieninger G. Fib420: a normal human variant of fibrinogen with two extended alpha chains. *Proc Natl Acad Sci U S A* 1994; 91(7):2625-2628.

- (72) Grieninger G, Lu X, Cao Y, Fu Y, Kudryk BJ, Galanakis DK et al. Fib420, the novel fibrinogen subclass: newborn levels are higher than adult. *Blood* 1997; 90(7):2609-2614.
- (73) Lishko VK, Yakubenko VP, Hertzberg KM, Grieninger G, Ugarova TP. The alternatively spliced alpha(E)C domain of human fibrinogen-420 is a novel ligand for leukocyte integrins alpha(M)beta(2) and alpha(X)beta(2). *Blood* 2001; 98(8):2448-2455.
- (74) Mosesson MW, Finlayson JS, Umfleet RA. Human fibrinogen heterogeneities. 3. Identification of chain variants. *J Biol Chem* 1972; 247(16):5223-5227.
- (75) Mosesson MW. Fibrinogen heterogeneity. *Ann N Y Acad Sci* 1983; 408:97-113.
- (76) Wolfenstein-Todel C, Mosesson MW. Human plasma fibrinogen heterogeneity: evidence for an extended carboxyl-terminal sequence in a normal gamma chain variant (gamma'). *Proc Natl Acad Sci U S A* 1980; 77(9):5069-5073.
- (77) Chung DW, Davie EW. gamma and gamma' chains of human fibrinogen are produced by alternative mRNA processing. *Biochemistry* 1984; 23(18):4232-4236.
- (78) Siebenlist KR, Meh DA, Mosesson MW. Plasma factor XIII binds specifically to fibrinogen molecules containing gamma chains. *Biochemistry* 1996; 35(32):10448-10453.
- (79) Kirschbaum NE, Mosesson MW, Amrani DL. Characterization of the gamma chain platelet binding site on fibrinogen fragment D. *Blood* 1992; 79(10):2643-2648.
- (80) Harfenist EJ, Packham MA, Mustard JF. Effects of variant gamma chains and sialic acid content of fibrinogen upon its interactions with ADP-stimulated human and rabbit platelets. *Blood* 1984; 64(6):1163-1168.
- (81) Bailey K, Bettelheim FR, Lorand L, Middlebrook WR. Action of thrombin in the clotting of fibrinogen. *Nature* 1951; 167(4241):233-234.
- (82) Blombäck B. Studies on the action of thrombotic enzymes on bovine fibrinogen as measured by N-terminal analysis. *Arkiv Kemi* 1958; 12:321-335.
- (83) Laurent T, Blombäck B. On the significance of the release of two different peptides from fibrinogen during clotting. *Acta Chemica Scandinavia* 1958; 12:1875-1877.
- (84) Magnusson S, Sottrup-Jensen L, Claeys H, Zajdel M, Petersen TE. Proceedings: Complete primary structure of prothrombin. Partial primary structures of plasminogen and hirudin. *Thromb Diath Haemorrh* 1975; 34(2):562-563.

- (85) Krishnaswamy S, Church WR, Nesheim ME, Mann KG. Activation of human prothrombin by human prothrombinase. Influence of factor Va on the reaction mechanism. *J Biol Chem* 1987; 262(7):3291-3299.
- (86) Tollefsen DM, Majerus DW, Blank MK. Heparin cofactor II. Purification and properties of a heparin-dependent inhibitor of thrombin in human plasma. *J Biol Chem* 1982; 257(5):2162-2169.
- (87) Rosenberg RD, Damus PS. The purification and mechanism of action of human antithrombin-heparin cofactor. *J Biol Chem* 1973; 248(18):6490-6505.
- (88) Pineda AO, Carrell CJ, Bush LA, Prasad S, Caccia S, Chen ZW et al. Molecular dissection of Na<sup>+</sup> binding to thrombin. *J Biol Chem* 2004; 279(30):31842-31853.
- (89) Di Cera E, Guinto ER, Vindigni A, Dang QD, Ayala YM, Wuyi M et al. The Na<sup>+</sup> binding site of thrombin. *J Biol Chem* 1995; 270(38):22089-22092.
- (90) Bale MD, Janmey PA, Ferry JD. Kinetics of formation of fibrin oligomers. II. Size distributions of ligated oligomers. *Biopolymers* 1982; 21(11):2265-2277.
- (91) Bailey K, Bettelheim F.R. The nature of the fibrinogen-thrombin reaction. *Br Med Bull* 1955; 11(1):50-53.
- (92) Martinelli RA, Scheraga HA. Steady-state kinetic study of the bovine thrombin-fibrinogen interaction. *Biochemistry* 1980; 19(11):2343-2350.
- (93) Hanna LS, Scheraga HA, Francis CW, Marder VJ. Comparison of structures of various human fibrinogens and a derivative thereof by a study of the kinetics of release of fibrinopeptides. *Biochemistry* 1984; 23(20):4681-4687.
- (94) Medved LV, Litvinovich SV, Ugarova TP, Lukinova NI, Kalikhevich VN, Ardemasova ZA. Localization of a fibrin polymerization site complementary to Gly-His-Arg sequence. *FEBS Lett* 1993; 320(3):239-242.
- (95) Shainoff JR, Dardik BN. Fibrinopeptide B in fibrin assembly and metabolism: physiologic significance in delayed release of the peptide. *Ann N Y Acad Sci* 1983; 408:254-268.
- (96) Hantgan RR, Hermans J. Assembly of fibrin. A light scattering study. *J Biol Chem* 1979; 254(22):11272-11281.
- (97) Jakobsen E, Ly B, Kierulf P. Incorporation of fibrinogen into soluble fibrin complexes. *Thromb Res* 1974; 4(4):499-507.
- (98) Harder AJ, Straub PW. In vitro and in vivo induction of cryofibrinogen and "paracoagulation" by reptilase. *Thromb Diath Haemorrh* 1972; 27(2):337-348.

- (99) Blomback B, Carlsson K, Fatah K, Hessel B, Procyk R. Fibrin in human plasma: gel architectures governed by rate and nature of fibrinogen activation. *Thromb Res* 1994; 75(5):521-538.
- (100) Blomback B, Banerjee D, Carlsson K, Hamsten A, Hessel B, Procyk R et al. Native fibrin gel networks and factors influencing their formation in health and disease. *Adv Exp Med Biol* 1990; 281:1-23.
- (101) Scott EM, Ariens RA, Grant PJ. Genetic and environmental determinants of fibrin structure and function: relevance to clinical disease. *Arterioscler Thromb Vasc Biol* 2004; 24(9):1558-1566.
- (102) Greenberg CS, Miraglia CC, Rickles FR, Shuman MA. Cleavage of blood coagulation factor XIII and fibrinogen by thrombin during in vitro clotting. *J Clin Invest* 1985; 75(5):1463-1470.
- (103) Chen R, Doolittle RF. Identification of the polypeptide chains involved in the cross-linking of fibrin. *Proc Natl Acad Sci U S A* 1969; 63(2):420-427.
- (104) Chen R, Doolittle RF. Cross-linking sites in human and bovine fibrin. *Biochemistry* 1971; 10(24):4487-4491.
- (105) McKee PA, Mattock P, Hill RL. Subunit structure of human fibrinogen, soluble fibrin, and cross-linked insoluble fibrin. *Proc Natl Acad Sci U S A* 1970; 66(3):738-744.
- (106) Nossel HL. Radioimmunoassay of fibrinopeptides in relation to intravascular coagulation and thrombosis. *N Engl J Med* 1976; 295(8):428-432.
- (107) Mills D, Karparkin S. The non-plasmin, proteolytic origin of human fibrinogen heterogeneity. *Biochim Biophys Acta* 1971; 251(1):121-125.
- (108) Hasegawa H, Nagata H, Yamauchi M, Murakoshi T, Oguma Y, Takei H. Synthesis, metabolism and distribution of fibrinogen and its derivatives. *Nippon Ketsueki Gakkai Zasshi* 1981; 44(7):1488-1498.
- (109) Holm B, Nilsen DW, Godal HC. Evidence that low molecular fibrinogen (LMW) is formed in man by degradation of high molecular weight fibrinogen (HMW). *Thromb Res* 1986; 41(6):879-884.
- (110) Sherman LA. Fibrinogen turnover: demonstration of multiple pathways of catabolism. *J Lab Clin Med* 1972; 79(5):710-723.
- (111) Pizzo SV, Schwartz ML, Hill RL, McKee PA. The effect of plasmin on the subunit structure of human fibrinogen. *J Biol Chem* 1972; 247(3):636-645.



- (112) Marder VJ, Shulman NR. High molecular weight derivatives of human fibrinogen produced by plasmin. II. Mechanism of their anticoagulant activity. *J Biol Chem* 1969; 244(8):2120-2124.
- (113) Gaffney PJ, Brasher M. Subunit structure of the plasmin-induced degradation products of crosslinked fibrin. *Biochim Biophys Acta* 1973; 295(1):308-313.
- (114) Pizzo SV, Taylor LM, Jr., Schwartz ML, Hill RL, McKee PA. Subunit structure of fragment D from fibrinogen and cross-linked fibrin. *J Biol Chem* 1973; 248(13):4584-4590.
- (115) Gaffney PJ. Distinction between fibrinogen and fibrin degradation products in plasma. *Clin Chim Acta* 1975; 65(1):109-115.
- (116) Bach-Gansmo ET, Halvorsen S, Godal HC, Skjongsberg OH. D-dimers are degraded by human neutrophil elastase. *Thromb Res* 1996; 82(2):177-186.
- (117) Wilhelmsen L, Svarsdudd K, Korsan-Bengtson K, Larsson B, Welin L, Tibblin G. Fibrinogen as a risk factor for stroke and myocardial infarction. *N Engl J Med* 1984; 311(8):501-505.
- (118) Baker IA, Pickering J, Elwood PC, Bayer A, Ebrahim S. Fibrinogen, viscosity and white blood cell count predict myocardial, but not cerebral infarction: evidence from the Caerphilly and Speedwell cohort. *Thromb Haemost* 2002; 87(3):421-425.
- (119) Ernst E, Resch KL. Fibrinogen as a cardiovascular risk factor: a meta-analysis and review of the literature. *Ann Intern Med* 1993; 118(12):956-963.
- (120) Meade TW, Ruddock V, Stirling Y, Chakrabarti R, Miller GJ. Fibrinolytic activity, clotting factors, and long-term incidence of ischaemic heart disease in the Northwick Park Heart Study. *Lancet* 1993; 342(8879):1076-1079.
- (121) Danesh J, Collins R, Appleby P, Peto R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA* 1998; 279(18):1477-1482.
- (122) Haines AP, Howarth D, North WR, Goldenberg E, Stirling Y, Meade TW et al. Haemostatic variables and the outcome of myocardial infarction. *Thromb Haemost* 1983; 50(4):800-803.
- (123) Reganon E, Vila V, Ferrando F, Martinez-Sales V, Fayos L, Ruano M et al. Elevated high molecular weight fibrinogen in plasma is predictive of coronary ischemic events after acute myocardial infarction. *Thromb Haemost* 1999; 82(5):1403-1405.

- (124) Qizilbash N, Jones L, Warlow C, Mann J. Fibrinogen and lipid concentrations as risk factors for transient ischaemic attacks and minor ischaemic strokes. *BMJ* 1991; 303(6803):605-609.
- (125) Ross R. Atherosclerosis--an inflammatory disease. *N Engl J Med* 1999; 340(2):115-126.
- (126) Lane DW, l'Anson S. Viscosimetric effect of fibrinogen. *J Clin Pathol* 1994; 47(11):1004-1005.
- (127) Blattler W, Straub PW, Jeanneret C, Horak GS. Effect of low fibrinogen concentrations on the rheology of human blood in vitro. *Am J Physiol* 1979; 236(3):H447-H450.
- (128) Merrill EW, Cokelet GC, Britten A, Wells REJr. Non-newtonian rheology of human blood--effect of fibrinogen deduced by "subtraction". *Circ Res* 1963; 13:48-55.
- (129) Chooi CC, Gallus AS. Acute phase reaction, fibrinogen level and thrombus size. *Thromb Res* 1989; 53(5):493-501.
- (130) Xiao Q, Danton MJ, Witte DP, Kowala MC, Valentine MT, Degen JL. Fibrinogen deficiency is compatible with the development of atherosclerosis in mice. *J Clin Invest* 1998; 101(5):1184-1194.
- (131) Lowe GD. Blood rheology in arterial disease. *Clin Sci* 1986; 71(2):137-146.
- (132) Rosencranz R, Bogen SA. Clinical laboratory measurement of serum, plasma, and blood viscosity. *American Journal of Clinical Pathology* 125 Suppl:S78-86, 2006.
- (133) Harkness J. The viscosity of human blood plasma; its measurement in health and disease. *Biorheology* 1971; 8(3):171-193.
- (134) Carroll S, Cooke CB, Butterly RJ. Plasma viscosity and its biochemical predictors: associations with lifestyle factors in healthy middle-aged men. *Blood Coagul Fibrinolysis* 2000; 11(7):609-616.
- (135) Blann A, Bignell A, McCollum C. von Willebrand factor, fibrinogen and other plasma proteins as determinants of plasma viscosity. *Atherosclerosis* 1998; 139(2):317-322.
- (136) Medzhitov R, Janeway CA, Jr. Innate immunity: impact on the adaptive immune response. *Curr Opin Immunol* 1997; 9(1):4-9.
- (137) Suffredini AF, Fantuzzi G, Badolato R, Oppenheim JJ, O'Grady NP. New insights into the biology of the acute phase response. *J Clin Immunol* 1999; 19(4):203-214.

- (138) Kopf M, Baumann H, Freer G, Freudenberg M, Lamers M, Kishimoto T et al. Impaired immune and acute-phase responses in interleukin-6-deficient mice. *Nature* 1994; 368(6469):339-342.
- (139) Benigni F, Fantuzzi G, Sacco S, Sironi M, Pozzi P, Dinarello CA et al. Six different cytokines that share GP130 as a receptor subunit, induce serum amyloid A and potentiate the induction of interleukin-6 and the activation of the hypothalamus-pituitary-adrenal axis by interleukin-1. *Blood* 1996; 87(5):1851-1854.
- (140) Baumann H, Gauldie J. The acute phase response. *Immunol Today* 1994; 15(2):74-80.
- (141) Seydewitz HH, Witt I. Increased phosphorylation of human fibrinopeptide A under acute phase conditions. *Thromb Res* 1985; 40(1):29-39.
- (142) Forsgren M, Raden B, Israelsson M, Larsson K, Heden LO. Molecular cloning and characterization of a full-length cDNA clone for human plasminogen. *FEBS Lett* 1987; 213(2):254-260.
- (143) Wallen P, Wiman B. Characterization of human plasminogen. II. Separation and partial characterization of different molecular forms of human plasminogen. *Biochim Biophys Acta* 1972; 257(1):122-134.
- (144) Robbins KC, Summari L, Hsieh B, Shah RJ. The peptide chains of human plasmin. Mechanism of activation of human plasminogen to plasmin. *J Biol Chem* 1967; 242(10):2333-2342.
- (145) Saksela O. Plasminogen activation and regulation of pericellular proteolysis. *Biochim Biophys Acta* 1985; 823(1):35-65.
- (146) Albrechtson OK. The fibrinolytic activity of human tissues. *Br J Haematol* 1957; 3(3):284-291.
- (147) Dano K, Andreasen PA, Grondahl-Hansen J, Kristensen P, Nielsen LS, Skriver L. Plasminogen activators, tissue degradation, and cancer. *Adv Cancer Res* 1985; 44:139-266.
- (148) Pennica D, Holmes WE, Kohr WJ, Harkins RN, Vehar GA, Ward CA et al. Cloning and expression of human tissue-type plasminogen activator cDNA in *E. coli*. *Nature* 1983; 301(5897):214-221.
- (149) Thorsen S. The mechanism of plasminogen activation and the variability of the fibrin effector during tissue-type plasminogen activator-mediated fibrinolysis. *Ann N Y Acad Sci* 1992; 667:52-63.

- (150) Halvorsen S, Skjonsberg OH, Godal HC. The stimulatory capacity of soluble fibrin prepared from high and low molecular weight fibrinogen on plasminogen activation. *Blood Coagul Fibrinolysis* 1993; 4(1):133-137.
- (151) Hoylaerts M, Rijken DC, Lijnen HR, Collen D. Kinetics of the activation of plasminogen by human tissue plasminogen activator. Role of fibrin. *J Biol Chem* 1982; 257(6):2912-2919.
- (152) Collen D. On the regulation and control of fibrinolysis. Edward Kowalski Memorial Lecture. *Thromb Haemost* 1980; 43(2):77-89.
- (153) Gurewich V, Pannell R, Louie S, Kelley P, Suddith RL, Greenlee R. Effective and fibrin-specific clot lysis by a zymogen precursor form of urokinase (pro-urokinase). A study in vitro and in two animal species. *J Clin Invest* 1984; 73(6):1731-1739.
- (154) Lijnen HR, Van Hoef B, Nelles L, Collen D. Plasminogen activation with single-chain urokinase-type plasminogen activator (scu-PA). Studies with active site mutagenized plasminogen (Ser740-Ala) and plasmin-resistant scu-PA (Lys158-Glu). *J Biol Chem* 1990; 265(9):5232-5236.
- (155) Ellis V, Behrendt N, Dano K. Plasminogen activation by receptor-bound urokinase. A kinetic study with both cell-associated and isolated receptor. *J Biol Chem* 1991; 266(19):12752-12758.
- (156) Moroi M, Aoki N. Isolation and characterization of alpha2-plasmin inhibitor from human plasma. A novel proteinase inhibitor which inhibits activator-induced clot lysis. *J Biol Chem* 1976; 251(19):5956-5965.
- (157) Bangert K, Johnsen AH, Christensen U, Thorsen S. Different N-terminal forms of alpha 2-plasmin inhibitor in human plasma. *Biochem J* 1993; 291 ( Pt 2):623-625.
- (158) Wiman B, Collen D. On the mechanism of the reaction between human alpha 2-antiplasmin and plasmin. *J Biol Chem* 1979; 254(18):9291-9297.
- (159) Aoki N, Harpel PC. Inhibitors of the fibrinolytic enzyme system. *Semin Thromb Hemost* 1984; 10:24-41.
- (160) Ny T, Sawdey M, Lawrence D, Millan JL, Loskutoff DJ. Cloning and sequence of a cDNA coding for the human beta-migrating endothelial-cell-type plasminogen activator inhibitor. *Proc Natl Acad Sci U S A* 1986; 83(18):6776-6780.
- (161) Pannekoek H, Veerman H, Lambers H, Diergaarde P, Verweij CL, van Zonneveld AJ et al. Endothelial plasminogen activator inhibitor (PAI): a new member of the Serpin gene family. *EMBO J* 1986; 5(10):2539-2544.

- (162) Lijnen HR. Pleiotropic functions of plasminogen activator inhibitor-1. *J Thromb Haemost* 2005; 3(1):35-45.
- (163) Wang W, Boffa MB, Bajzar L, Walker JB, Nesheim ME. A study of the mechanism of inhibition of fibrinolysis by activated thrombin-activable fibrinolysis inhibitor. *J Biol Chem* 1998; 273(42):27176-27181.
- (164) Redlitz A, Tan AK, Eaton DL, Plow EF. Plasma carboxypeptidases as regulators of the plasminogen system. *J Clin Invest* 1995; 96(5):2534-2538.
- (165) Jacobsson K. Studies on the determination of fibrinogen in human blood plasma. *Scand J Clin Lab Invest* 1955; 7(Suppl. 14):1-54.
- (166) Clauss A. Rapid physiological coagulation method in determination of fibrinogen. *Acta Haematol* 1957; 17(4):237-246.
- (167) Desvignes P, Bonnet P. Direct determination of plasma fibrinogen levels by heat precipitation. A comparison of the technique against thrombin clottable fibrinogen with spectrophotometry and radial immuno-diffusion. *Clin Chim Acta* 1981; 110(1):9-17.
- (168) Exner T, Burrige J, Power P, Rickard KA. An evaluation of currently available methods for plasma fibrinogen. *Am J Clin Pathol* 1979; 71(5):521-527.
- (169) Rampling MW, Gaffney PJ. The sulphite precipitation method for fibrinogen measurement; its use on small samples in the presence of fibrinogen degradation products. *Clin Chim Acta* 1976; 67(1):43-52.
- (170) Hoegge-de Nobel E, Voskuilen M, Briet E, Brommer EJ, Nieuwenhuizen W. A monoclonal antibody-based quantitative enzyme immunoassay for the determination of plasma fibrinogen concentrations. *Thromb Haemost* 1988; 60(3):415-418.
- (171) Palareti G, Maccaferri M, Manotti C, Tripodi A, Chantarangkul V, Rodeghiero F et al. Fibrinogen assays: a collaborative study of six different methods. C.I.S.M.E.L. Comitato Italiano per la Standardizzazione dei Metodi in Ematologia e Laboratorio. *Clin Chem* 1991; 37(5):714-719.
- (172) Gaffney PJ, Wong MY. Collaborative study of a proposed international standard for plasma fibrinogen measurement. *Thromb Haemost* 1992; 68(4):428-432.
- (173) Whitton CM, Sands D, Hubbard AR, Gaffney PJ. A collaborative study to establish the 2nd International Standard for Fibrinogen, Plasma. *Thromb Haemost* 2000; 84(2):258-262.
- (174) Arnaout MA. Structure and function of the leukocyte adhesion molecules CD11/CD18. *Blood* 1990; 75(5):1037-1050.

- (175) Weiss HJ, Hawiger J, Ruggeri ZM, Turitto VT, Thiagarajan P, Hoffmann T. Fibrinogen-independent platelet adhesion and thrombus formation on subendothelium mediated by glycoprotein IIb-IIIa complex at high shear rate. *J Clin Invest* 1989; 83(1):288-297.
- (176) Ikeda Y, Handa M, Kawano K, Kamata T, Murata M, Araki Y et al. The role of von Willebrand factor and fibrinogen in platelet aggregation under varying shear stress. *J Clin Invest* 1991; 87(4):1234-1240.
- (177) Bennett JS. Platelet-fibrinogen interactions. *Ann N Y Acad Sci* 2001; 936:340-354.
- (178) Farrell DH, Thiagarajan P, Chung DW, Davie EW. Role of fibrinogen alpha and gamma chain sites in platelet aggregation. *Proc Natl Acad Sci U S A* 1992; 89(22):10729-10732.
- (179) Flick MJ, Du X, Witte DP, Jirouskova M, Soloviev DA, Busuttill SJ et al. Leukocyte engagement of fibrin(ogen) via the integrin receptor alphaMbeta2/Mac-1 is critical for host inflammatory response in vivo. *J Clin Invest* 2004; 113(11):1596-1606.
- (180) Altieri DC, Plescia J, Plow EF. The structural motif glycine 190-valine 202 of the fibrinogen gamma chain interacts with CD11b/CD18 integrin (alpha M beta 2, Mac-1) and promotes leukocyte adhesion. *J Biol Chem* 1993; 268(3):1847-1853.
- (181) Altieri DC, Mannucci PM, Capitanio AM. Binding of fibrinogen to human monocytes. *J Clin Invest* 1986; 78(4):968-976.
- (182) Fan ST, Edgington TS. Integrin regulation of leukocyte inflammatory functions. CD11b/CD18 enhancement of the tumor necrosis factor-alpha responses of monocytes. *J Immunol* 1993; 150(7):2972-2980.
- (183) Pakianathan DR. Extracellular matrix proteins and leukocyte function. *J Leukoc Biol* 1995; 57(5):699-702.
- (184) Perez RL, Roman J. Fibrin enhances the expression of IL-1 beta by human peripheral blood mononuclear cells. Implications in pulmonary inflammation. *J Immunol* 1995; 154(4):1879-1887.
- (185) Doherty DE, Haslet C, Tonnesen MG, Henson PM. Human monocyte adherence : a primary effect of chemotactic factors on the monocyte to stimulate adherence to human endothelium. *J Immunol* 1987; 138:1762-1771.
- (186) Ciano PS, Colvin AM, Dvorak AM. Macrophage migration in fibrin gel matrices. *Lab Invest* 1986; 54:62-70.

- (187) Tang L, Eaton JW. Fibrin(ogen) mediates acute inflammatory responses to biomaterials. *J Exp Med* 1993; 178(6):2147-2156.
- (188) Tang L, Ugarova TP, Plow EF, Eaton JW. Molecular determinants of acute inflammatory responses to biomaterials. *J Clin Invest* 1996; 97(5):1329-1334.
- (189) Cheresch DA, Berliner SA, Vicente V, Ruggeri AM. Recognition of distinct adhesive sites on fibrinogen by related integrins on platelets and endothelial cells. *Cell* 1989; 58:945-953.
- (190) Suehiro K, Gailit J, Plow EF. Fibrinogen is a ligand for integrin  $\alpha 5 \beta 1$  on endothelial cells. *J Biol Chem* 1997; 272:5360-5366.
- (191) Clauss A. [Rapid physiological coagulation method in determination of fibrinogen.]. *Acta Haematol* 1957; 17(4):237-246.
- (192) Campbell PJ. International biological standards and reference preparations. I. Preparation and presentation of materials to serve as standards and reference preparations. *J Biol Stand* 1974; 2(4):249-258.
- (193) Weinstein MJ, Deykin D. Quantitative abnormality of an Aalpha chain molecular weight form in the fibrinogen of cirrhotic patients. *Br J Haematol* 1978; 40(4):617-630.
- (194) Jakobsen K, Kierulf P. A modified beta-alanine precipitation procedure to prepare fibrinogen free of anti-thrombin III and plasminogen. *Thromb Res* 1973; 3:145-159.
- (195) Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25(4):402-408.
- (196) Gaffney PJ, Wong MY. Collaborative study of a proposed international standard for plasma fibrinogen measurement. *Thromb Haemost* 1992; 68(4):428-432.
- (197) Furlan M, Felix R, Escher N, Lammle B. How high is the true fibrinogen content of fibrinogen standards? *Thromb Res* 1989; 56(5):583-592.
- (198) Schwegman JJ, Carpenter JF, Nail SL. Infrared microscopy for in situ measurement of protein secondary structure during freezing and freeze-drying. *J Pharm Sci* 2007; 96(1):179-195.
- (199) Prestrelski SJ, Arakawa T, Carpenter JF. Separation of freezing- and drying-induced denaturation of lyophilized proteins using stress-specific stabilization. II. Structural studies using infrared spectroscopy. *Arch Biochem Biophys* 1993; 303(2):465-473.

- (200) Ly B, Godal HC, Kierulf P. Thrombin clotting time--a more sensitive indicator of denatured fibrinogen than clottability. *Haemostasis* 1972; 1(1):44-51.
- (201) Holm B, Godal HC. Quantitation of the three normally-occurring plasma fibrinogens in health and during so-called "acute phase" by SDS electrophoresis of fibrin obtained from EDTA-plasma. *Thromb Res* 1984; 35(3):279-290.
- (202) Holm B, Nilsen DW, Kierulf P, Godal HC. Purification and characterization of 3 fibrinogens with different molecular weights obtained from normal human plasma. *Thromb Res* 1985; 37(1):165-176.
- (203) Blomback B, Blomback M, Laurent TC, Pertoft H. Effect of EDTA on fibrinogen. *Biochim Biophys Acta* 1966; 127(2):560-562.
- (204) Godal HC. The effect of EDTA on human fibrinogen and its significance for the coagulation with thrombin. Oslo University Press 1960;6-7.
- (205) Joly M. Recent studies of the reversible denaturation of proteins. *Prog Biophys* 1955; 5:168-220.
- (206) Keil U, Chambless LE, Doring A, Filipiak B, Stieber J. The relation of alcohol intake to coronary heart disease and all-cause mortality in a beer-drinking population. *Epidemiology* 1997; 8(2):150-156.
- (207) Klatsky AL, Friedman GD, Siegelaub AB. Alcohol use and cardiovascular disease: the Kaiser-Permanente experience. *Circulation* 1981; 64(3 Pt 2):III-41.
- (208) Rimm EB, Giovannucci EL, Willett WC, Colditz GA, Ascherio A, Rosner B et al. Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet* 1991; 338(8765):464-468.
- (209) Retterstol L, Berge KE, Braaten O, Eikvar L, Pedersen TR, Sandvik L. A daily glass of red wine: does it affect markers of inflammation? *Alcohol Alcohol* 2005; 40(2):102-105.
- (210) Lowe GD, Drummond MM, Forbes CD, Barbenel JC. The effects of age and cigarette-smoking on blood and plasma viscosity in men. *Scott Med J* 1980; 25(1):13-17.
- (211) Harkness J. The viscosity of human blood plasma; its measurement in health and disease. *Biorheology* 1971; 8(3):171-193.
- (212) Otto C, Richter WO, Schwandt P. Contribution of fibrinogen and lipoproteins to plasma viscosity in hypercholesterolemia and hypertriglyceridemia: evaluation by selective depletion of low-density lipoproteins or fibrinogen. *Metabolism* 2000; 49(6):810-813.



- (213) Rand PW, Lacombe E, Hunt HE, Austin WH. Viscosity of normal human blood under normothermic and hypothermic conditions. *J Appl Physiol* 1964; 19:117-122.
- (214) Woodward M, Lowe G, Rumley A, Imhof A, Koenig W. Measurement of plasma viscosity in stored frozen samples: a general population study. *Blood Coagul Fibrinolysis* 2003; 14(4):417-420.
- (215) Mukamal KJ, Jadhav PP, D'Agostino RB, Massaro JM, Mittleman MA, Lipinska I et al. Alcohol consumption and hemostatic factors: analysis of the Framingham Offspring cohort. *Circulation* 2001; 104(12):1367-1373.
- (216) Lowe GD, Wood DA, Douglas JT, Riemersma RA, Macintyre CC, Takase T et al. Relationships of plasma viscosity, coagulation and fibrinolysis to coronary risk factors and angina. *Thromb Haemost* 1991; 65(4):339-343.
- (217) Sweetnam PM, Thomas HF, Yarnell JW, Beswick AD, Baker IA, Elwood PC. Fibrinogen, viscosity and the 10-year incidence of ischaemic heart disease. *Eur Heart J* 1996; 17(12):1814-1820.
- (218) Levi M, Van Der PT. Two-way interactions between inflammation and coagulation. *Trends Cardiovasc Med* 2005; 15(7):254-259.
- (219) Van Der PT, de Jonge E, Levi M. Regulatory role of cytokines in disseminated intravascular coagulation. *Semin Thromb Hemost* 2001; 27(6):639-651.
- (220) Fan ST, Edgington TS. Coupling of the adhesive receptor CD11b/CD18 to functional enhancement of effector macrophage tissue factor response. *J Clin Invest* 1991; 87(1):50-57.
- (221) Labrador V, Legrand S, Muller S, Carl P, Senger B, Voegel J-C et al. Adhesion of leukocytes on a fibrinogen-coated surface in static conditions: Experimentation and modelization. [French]. *Journal des Maladies Vasculaires* Vol 25(1)(pp 47-52), 2000 2000;(1):47-52.
- (222) Languino LR, Plescia J, Duperray A, Brian AA, Plow EF, Geltosky JE et al. Fibrinogen mediates leukocyte adhesion to vascular endothelium through an ICAM-1-dependent pathway. *Cell* 1993; 73(7):1423-1434.
- (223) Trezzini C, Schuepp B, Maly FE, Jungi TW. Evidence that exposure to fibrinogen or to antibodies directed against Mac-1 (CD11b/CD18; CR3) modulates human monocyte effector functions. *Br J Haematol* 1991; 77(1):16-24.
- (224) Smiley ST, King JA, Hancock WW. Fibrinogen Stimulates Macrophage Chemokine Secretion Through Toll-Like Receptor 4. *J Immunol* 2001; 167(5):2887-2894.

- (225) Ugarova TP, Yakubenko VP. Recognition of fibrinogen by leukocyte integrins. *Annals of the New York Academy of Sciences* 2001; 936:368-385.
- (226) Perez RL, Ritzenthaler JD, Roman J. Transcriptional regulation of the interleukin-1beta promoter via fibrinogen engagement of the CD18 integrin receptor. *Am J Respir Cell Mol Biol* 1999; 20(5):1059-1066.
- (227) Sitrin RG, Pan PM, Srikanth S, Todd RF, III. Fibrinogen activates NF-kappa B transcription factors in mononuclear phagocytes. *J Immunol* 1998; 161(3):1462-1470.
- (228) De Groote D, Zangerle PF, Gevaert Y, Fassotte MF, Beguin Y, Noizat-Pirenne F et al. Direct stimulation of cytokines (IL-1 beta, TNF-alpha, IL-6, IL-2, IFN-gamma and GM-CSF) in whole blood. I. Comparison with isolated PBMC stimulation. *Cytokine* 1992; 4(3):239-248.
- (229) Oliver JC, Bland LA, Oettinger CW, Arduino MJ, McAllister SK, Aguero SM et al. Cytokine kinetics in an in vitro whole blood model following an endotoxin challenge. *Lymphokine Cytokine Res* 1993; 12(2):115-120.
- (230) Barth S, Kleinhapp B, Gutschi A, Jelovcan S, Marth E. In vitro cytokine mRNA expression in normal human peripheral blood mononuclear cells. *Inflamm Res* 2000; 49(6):266-274.
- (231) DeForge LE, Remick DG. Kinetics of TNF, IL-6, and IL-8 gene expression in LPS-stimulated human whole blood. *Biochem Biophys Res Commun* 1991; 174(1):18-24.
- (232) Cavaillon JM, Fitting C, Haeffner-Cavaillon N, Kirsch SJ, Warren HS. Cytokine response by monocytes and macrophages to free and lipoprotein-bound lipopolysaccharide. *Infect Immun* 1990; 58(7):2375-2382.
- (233) Kirikae T, Tamura H, Hashizume M, Kirikae F, Uemura Y, Tanaka S et al. Endotoxin contamination in fetal bovine serum and its influence on tumor necrosis factor production by macrophage-like cells J774.1 cultured in the presence of the serum. *Int J Immunopharmacol* 1997; 19(5):255-262.
- (234) Busso N, Peclat V, Van Ness K, Kolodzieszyk E, Degen J, Bugge T et al. Exacerbation of antigen-induced arthritis in urokinase-deficient mice. *J Clin Invest* 1998; 102(1):41-50.
- (235) Dvorak HF, Senger DR, Dvorak AM, Harvey VS, McDonagh J. Regulation of extravascular coagulation by microvascular permeability. *Science* 1985; 227(4690):1059-1061.
- (236) Martin-Paredero V, Vadillo J, Diaz J, Espinosa A, Berga C, Segura J et al. Fibrinogen and fibrinolysis in blood and in the arterial wall: its role in advanced atherosclerotic disease. *Cardiovasc Surg* 1998; 6(5):457-462.

- (237) Bini A, Kudryk BJ. Fibrinogen and fibrin in the arterial wall. *Thromb Res* 1994; 75(3):337-341.